



E-ISSN: 2320-7078
P-ISSN: 2349-6800
JEZS 2017; 5(3): 176-179
© 2017 JEZS
Received: 26-03-2017
Accepted: 27-04-2017

Sharma P
Department of Agricultural
Entomology, S.K.N. College of
Agriculture, Jobner, Jaipur,
India

Kumawat KC
Department of Agricultural
Entomology, S.K.N. College of
Agriculture, Jobner, Jaipur,
India

Jhumar Lal
Department of Entomology,
RCA, MPUAT, Udaipur,
Rajasthan, India

Seasonal abundance of diamondback moth and natural enemies on cabbage

Sharma P, Kumawat KC and Jhumar Lal

Abstract

The cabbage crop was found to be infested by diamondback moth (*Plutella xylostella* L.). The infestation of diamondback moth started from the third week of November and reached peak (45.2 larvae /10 plants) in the first week of January. The maximum and minimum temperature showed significant negative correlation with larval population of diamondback moth whereas, non-significant correlation with relative humidity and sunshine hours. The coccinellid beetle was recorded as an important predator of aphid, which was maximum (20.2 /10 plants) in the fourth week of January.

Keywords: Diamondback moth, *Plutella xylostella*, coccinellid, cabbage, seasonal abundance and natural enemies

1. Introduction

Cabbage, *Brassica oleracea* var. *capitata* L. is one of the important cruciferous vegetable crops grown in India. It is grown more or less in all the states. The nutritional value/ 100 g of cabbage consists of carbohydrates 5.8 g, fat 0.1 g, protein 1.28 g, vitamins (thiamine or vitamin B₁ 0.061 mg, riboflavin or B₂ 0.040 mg, niacin or vitamin B₃ 0.234 mg, pantothenic acid or vitamin B₅ 0.212 mg, folate or vitamin B₉ 43 mg, vitamin C 36.6 mg, and vitamin K 76 mg), minerals (Ca 40 mg, Fe 0.47 mg, Mg 12 mg, Mn 0.16 mg, P 26 mg, K 170 mg, Na 18 mg, Zn 0.18 mg) [1]. The total area under cultivation of cabbage in India is 372 thousand hectares with an annual production to the tune of 8534 thousand tonnes with productivity of 18.3 metric tonnes [2]. The total area under cultivation of cabbage in Rajasthan is 346 hectares with an annual production to the tune of 7588 tonnes [3]. China is major cabbage producing country with 47 per cent of world followed by India with 12 per cent of world production [4]. The yield of cabbage is adversely affected by many bottlenecks including insect pest, diseases, environmental stresses, nutritional imbalance etc. Among them, insect pests, viz., tobacco caterpillar, *Spodoptera litura* (Fab.); diamondback moth, *Plutella xylostella* (L.) cabbage borer, *Hellula undalis* Fab.; cabbage looper, *Tricoplusia ni* Hub and aphid, *Lipaphis erysimi* (Kalt.) [5-9]. Out of these, aphid and diamondback moth are major pests causing significant loss in North India. The diamondback moth, *P. xylostella* was first reported on cruciferous vegetables in 1914 [10]. It is sometimes called cabbage moth. So, to have clear proven idea on the seasonal incidence of diamondback moth and natural enemies in cabbage ecosystem and their relation with abiotic factors like temperature and relative humidity, the present experiment was conducted for further understanding of the role played by the abiotic factors in the incidence of diamondback moth which is harmful to cabbage growers which ultimately will help cabbage growers for better return in terms of yield as well as income generation.

2. Materials and Methods

2.1 site and location of experiments: The present investigations were conducted at Horticulture Farm of S.K.N. College of Agriculture, Jobner, Jaipur during *Rabi*, 2015-16. Geographically, Jobner is located at longitude of 75°28' East, latitude of 26°06' North and at an altitude of 427 metres above mean sea level (MSL) in Jaipur district of Rajasthan.

2.2 Climatic and weather conditions of location: The climate of the region is typically semi-arid which is characterized by extremes of the temperature during summer and winter. During summer, temperature may rise as high as 47 °C and in winter, it may fall as low as 2-3 °C. The total rainfall is 500 mm which is mostly received from last week of June to September. This region provides a safe long growing season for most of the crops.

Correspondence
Jhumar Lal
Department of Entomology,
RCA, MPUAT, Udaipur,
Rajasthan, India

2.3 Layout and design of Experiment: The experiment was laid out in five plots of 2.25 x 2.25 m² sizes during *Rabi*, 2015-16. These plots were contiguous to each. Five weeks old seedlings were transplanted at row to row and plant to plant distance of 45 x 45 cm, respectively. The transplanting was done on 20th October during *Rabi*, 2015-16 in the evening hours followed by light irrigation. Other cultural practices as recommended in Packages of Practices were followed during the experimental period.

2.4 Observations: Observations on the abundance of diamondback moth and natural enemies were recorded from very beginning of their appearance on plants till harvesting of the crop.

2.4.1 Diamond back moth: In order to register the larval population of diamondback moth (DBM); direct visual counting method was used [11]. Ten plants were selected randomly from each plot and the total larval population of the pest was recorded at weekly interval.

2.4.2 Natural enemies: Coccinellid predator *C. septempunctata* was appeared as the major natural enemy in cabbage ecosystem; its population was recorded on ten randomly selected plants per plot.

2.4.3 Meteorological data: Weekly data of temperature (maximum and minimum), relative humidity and sunshine hours were collected from Meteorological observatory, Department of Agronomy, S.K.N. college of Agriculture, Jobner, Jaipur, and Rajasthan.

2.5 Statistical analysis: Population data of *Plutella xylostella* and natural enemy (Coccinellids) thus obtained were subjected to statistical analysis to find out the coefficient of correlation with average temperature relative humidity and sunshine hours. A simple correlation was worked out between the population of *P. xylostella* and natural enemy (Coccinellids) with abiotic environmental factor using the following formula.

$$r_{xy} = \frac{\sum XY - \frac{\sum X \sum Y}{n}}{\sqrt{\left[\sum X^2 - \frac{(\sum X)^2}{n} \right] \left[\sum Y^2 - \frac{(\sum Y)^2}{n} \right]}}$$

Where,

r_{xy} = Simple correlation coefficient
 X = Variable i.e. abiotic component. (Average temperature and relative humidity)
 Y = Variable i.e. mean number of insect pests
 n = Number of paired observations

$$t = \frac{r}{\sqrt{1 - r^2}} \times \sqrt{n - 2} \sim t_{n-2} \text{ d.f.}$$

Significance of correlation is tested by using following formula.

3. Results and Discussion

During the present study, the crop was found to be abundantly infested with diamondback moth, *P. xylostella*. Among the natural enemies of insect pests of cabbage, coccinellid predator, *Coccinella septempunctata* L. was found major in the cabbage ecosystem which feeds on the aphids. Other natural enemies like *Cotesia plutellae* and Syrphid flies were present in traces only. This insect pest has also been reported as serious insect pest of cabbage crop by some entomologists [6, 12, 13] who also support the present findings

3.1 Seasonal abundance of *P. xylostella*: The data is presented in Table 1 and revealed that the infestation of *P. xylostella* on cabbage was started from second week of November (47th SMW) and reached to maximum (45.2 larvae /10 plants) in the first week of January 1st SMW) during the year, thereafter, population started declining. The infestation starts from third week of November and reached to maximum (45.75 and 42.50 larvae/10 plants) in the first week of January thereafter; population started declined [14]. Peak population of diamond back moth (DBM) was recorded on 1st March and 23rd February with 13.60 and 14.33 larvae/plant during 2011-12 and 2012-13 respectively [15]. The peak population of diamondback moth was found in the first week of February [16]. The infestation of *P. xylostella* started from second week of November and reached to peak in the last week of January to first week of February [17-20]. The larval population of diamondback moth had significantly negative correlation with maximum ($r = -0.56$) and minimum temperatures ($r = -0.59$) and non-significant correlation with relative humidity. The studies showed that the incidence of diamondback moth was only affected by temperature, i.e. with rise in temperature, the population of pest declined. Population of diamondback moth was negatively correlated with mean temperature and non-significant with relative humidity [16]. Non-significant positive [13], and significant positive correlation [18], respectively were also recorded between the population of diamondback moth and temperature but these were controversy result. Population of diamondback moth started to build up after the cauliflower crop was transplanted [22].

3.2 Seasonal abundance of natural enemies *Coccinella septempunctata* L.: Among the natural enemies *Coccinella septempunctata* L., a predator of aphid was found in abundance. The data is presented in Table 1 and revealed that the population of *C. septempunctata* was maximum (20.2 /10 plants) in the fourth week of January (4th SMW) and had non-significant correlation with maximum and minimum temperatures, relative humidity and negative sunshine hours. The *C. septempunctata*, population was maximum (35.25 and 40.00/10 plants) in the third and second week of January, 2008-09 and 2009-10 [14], whereas maximum coccinellid was observed on 23rd February of 2011-12 and 2012-13 crop seasons with 11.67 and 9.67 coccinellids/ 5plants, respectively [15]. The population of *C. septempunctata* had non-significant correlation with maximum and minimum temperatures, relative humidity and negative sunshine hours [23, 24]. The results also get evidence from the finding of some entomologists that the parasitization of diamondback moth by *Cotesia plutellae* [25- 27] and *Tetrastichus sokolowskii* [28] were not recorded in cabbage ecosystem.

Table 1: Seasonal abundance of diamondback moth, *Plutella xylostella* (L.) and natural enemies in cabbage crop during rabi, 2015-16

S. No.	Date of observation	Standard meteorological week (SMW)	Temperature (°C)		Relative humidity (%)	Sunshine hours/day	<i>P. xylostella</i> Larvae / 10 plants	No. of <i>C. septempunctata</i> /10 plants
			Maximum	Minimum				
1	19/11/15	47	29.1	09.0	53	8.5	2.0	0.00
2	26/11/15	48	27.2	09.3	61	8.4	10.0	0.00
3	3/12/15	49	28.4	06.9	60	8.4	21.6	8.6
4	10/12/15	50	23.5	03.2	63	7.6	28.2	10.0
5	17/12/15	51	23.0	02.1	57	6.1	33.6	10.8
6	24/12/15	52	24.7	04.0	57	8.3	40.4	11.8
7	1/1/16	1	27.2	08.1	63	7.6	45.2	14.8
8	8/1/16	2	26.1	05.5	59	7.0	40.2	16.8
9	15/1/16	3	21.7	03.3	67	5.3	37.2	17.2
10	22/1/16	4	24.1	03.5	62	8.1	35.4	20.2
11	29/1/16	5	25.7	07.7	64	8.4	30.4	18.4
12	5/2/16	6	25.2	05.2	51	6.8	27.6	14.8
13	12/2/16	7	25.0	09.3	48	8.2	25.6	9.2
14	19/2/16	8	27.6	09.6	55	8.7	22.4	5.6
Seasonal Mean			25.61	6.19	58.57	7.67	28.56	11.30
Coefficient of correlation (r) for population and Maximum Temperature							-0.56*	-0.48
Coefficient of correlation (r) for population and Minimum Temperature							-0.55*	-0.37
Coefficient of correlation (r) for population and Relative Humidity							0.37	0.30
Coefficient of correlation (r) for population and Sunshine							-0.47	-0.37

* Significant at the 5 % level of significance

4. Conclusion

The seasonal abundance of diamondback moth of cabbage was studied that will be helpful in preparing proper schedule for effective management of this pest. During the present study, the crop was found to be abundantly infested with diamondback moth, *P. xylostella* throughout the growing season. Among the natural enemies of insect pests of cabbage, coccinellid predator, *Coccinella septempunctata* L. was appeared as the major natural enemy in cabbage ecosystem. Other natural enemies like *Cotesia* and Syrphid flies were presented in limited numbers. *C. septempunctata* was appeared as the major natural enemy in cabbage ecosystem.

5. References

- Anonymous, Nutritional value of different vegetable. Nutrition value. org. 2016.
- Anonymous. National Horticulture Board. www.nhb.com. 2013.
- Anonymous. Directorate of Horticulture, Government of Rajasthan, Jaipur, Rajasthan. 2010.
- Anonymous. Food and agriculture organization. FAO. Org. 2012.
- Prasad SK. Quantitative estimation of damage to cruciferous crops caused by cabbage worm, cabbage lopper, diamondback moth and cabbage aphid. Indian Journal of Entomology. 1963; 25(3):242-259.
- Sachan JN, Srivastava BP. Studies of the seasonal incidence of insect pests of cabbage. Indian Journal of Entomology. 1972; 34(2):123-129.
- Joshi FL, Sharma JC. Efficacy of different insecticidal treatment schedules against pests of cabbage. Labdev Journal of Science and Technology. Part B. 1976; 11:1-5.
- Mohan NJ, Krishnaiah K, Kumar NKK. Chemical control of mustard aphid, *Lipaphis erysimi* (Kalt). And leaf webber, *Crocidolomia binotalis* Zell. on cabbage. Pesticides. 1981; 15(2):29-32.
- Rao SRK, Lal OP. Seasonal incidence of mustard aphid, *Lipaphis erysimi* (Kalt) and diamondback moth, *Plutella xylostella* (L.) on cabbage. Journal of Insect Science. 2005; 18(2):106-110.
- Fletcher TB. Some south Indian insect. Superintendent govt. press, Madras, 1914, 565.
- Lal OP. Notes summer school on Advance technologies in Important Vegetable Crops, including Cole Crops. May 4-24, I.A.R.I. New Delhi. 1998, 63-66.
- Kandoria JL, Lal A, Singh L. Seasonal availability of diamondback moth, *Plutella xylostella* (L.) under Punjab conditions. Journal of Insect Science. 1996; 9(1):6-8.
- Sharma SK. Eco-safe management of major insect pests of cabbage Brassica oleracea var. capitata Linn. Ph.D Thesis, Submitted to Rajasthan Agricultural University, Bikaner, 2004.
- Bana JK, Jat BL, Bajya DR. Seasonal incidence of major insect pest of cabbage and their natural enemies. Indian Journal of Entomology. 2012; 74(3):236-240.
- Patra S, Dhote VW, Alam SKF, Das BC, Chatterjee ML, Samanta A. Population dynamics of major insect pests and their natural enemies on cabbage under new alluvial zone of West Bengal. The Journal of Plant Protection Sciences. 2013; 5(1):42-49.
- Khaire VA, Lauvande KE, Ajvi DS. Population dynamics of insect pests of cabbage. Current Research. 1987; 48(2):48-86.
- Gera SS, Bhatnagar KN. Seasonal incidence of pest complex of cabbage and their control in a semi-arid region. Pestology. 1992; 16(3):38-45.
- Meena SC, Singh V. Seasonal incidence of *Plutella xylostella* (L.) on cabbage in arid region of Rajasthan. Annals of Plant Protection Sciences. 2012; 20(3):326-328.
- Shukla A, Kumar A. Seasonal incidence of diamondback moth, *Plutella xylostella* (Linn.) in relation to abiotic factors in cabbage. Plant Protection Bulletin. 2004; 1:37-38.
- Goud CR, Rao SRK, Chiranjeevi CH. Influence of weather parameters on the population buildup of diamondback moth, *Plutella xylostella* (L.) infesting cabbage. Pest management in Horticulture Ecosystem. 2006; 12(1):103-106.
- Chaudhuri N, Ghosh S, Ghosh J, Senapati SK. Incidence of insect pests of cabbage in relation to prevailing climatic conditions of Terai region. Indian Journal of

- Entomology. 2001; 63:421-8.
22. Ahmad T, Ansari M. Studies on seasonal abundance of diamondback moth, *Plutella xylostella* on cauliflower crop. Journal of Plant Protection Research. 2010; 50(3):280-287.
 23. Chandra S, Kushawa KS. Impact of environmental resistance on aphid complex of cruciferous crops under the agro-climatic conditions of Udaipur II biotic components. Indian Journal of Entomology. 1987; 49(1):86-113.
 24. Mandal SMA, Patnaik NC. Predatory potential of aphidophagous predators associated with cabbage crop. Journal of Plant Protection Environment. 2006; 3(1):81-86.
 25. Shukla A, Kumar A. Seasonal activity of larval parasitoid, *Cotesia plutellae* (Kurdj.) in cabbage ecosystem. Journal of Entomological Research. 2003; 27(3):181-184.
 26. Bhaskar H, Viraktamath CA. Diversity and abundance of aphidophagous coccinellids in cabbage field. Insect Environment. 2002; 8(1):31.
 27. Bar UK, Uma Shankar, Raju SVS. Seasonal abundance and parasitization of diamondback moth and larval parasitoid, *Cotesia plutellae*. Annals of Plant Protection Science. 2001; 12(2):214-216.
 28. Shukla A, Kumar A. Seasonal activity of *Tetrastichus sokolowski* a pupal parasitoid of *Plutella xylostella* (Linn.) in cabbage ecosystem. Journal of Entomological Research. 2006; 30(2):155-157.