



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2018; 6(3): 142-145

© 2018 JEZS

Received: 08-03-2018

Accepted: 09-04-2018

Akansha SinghPh.D. Scholar, ICAR- Indian
Veterinary Research Institute,
Izatnagar, Bareilly, Uttar
Pradesh, India**Adesh Kumar**Veterinary Assistant Surgeon,
Madhya Pradesh Government**Babu Lal Saini**Ph.D. Scholar, ICAR- Indian
Veterinary Research Institute,
Izatnagar, Bareilly, Uttar
Pradesh, India**Arnav Mehrotra**Ph.D. Scholar, ICAR- Indian
Veterinary Research Institute,
Izatnagar, Bareilly, Uttar
Pradesh, India**Sandeep Kumar Chaudhary**Ph.D. Scholar, ICAR- Indian
Veterinary Research Institute,
Izatnagar, Bareilly, Uttar
Pradesh, India**Vandana Yadav**Ph.D. Scholar, ICAR- National
Dairy Research Institute,
Karnal, Haryana, India**Correspondence****Akansha Singh**Ph.D. Scholar, ICAR- Indian
Veterinary Research Institute,
Izatnagar, Bareilly, Uttar
Pradesh, India

Effect of non-genetic factors on milk fat composition: A review

Akansha Singh, Adesh Kumar, Babu Lal Saini, Arnav Mehrotra, Sandeep Kumar Chaudhary and Vandana Yadav

Abstract

Milk fat comprises of mono- and polyunsaturated fatty acids; some of them are recognized to exert a beneficial effect on human health. Short and medium chain saturated fatty acids are synthesized *de novo* in the mammary gland while long chain fatty acid are derived from blood circulation. Milk fat is the main source of energy and natural carriers' of triacylglycerols and bioactive molecules to the intestine of infants. They are also involved in providing the information needed for the growth of intestinal mucosa, the immune and nervous systems along with metabolic activity. They have impact on the technological and nutritional properties of the milk, hence, are important components of milk composition. The composition of milk fatty acid is affected by numerous non genetic factors such as species differences, parity or lactation order, age, size and health status of the cow, dietary composition, feeding, season, pregnancy, sire service period, lameness, heat stress, milking interval *etc.* Changes in milk fatty acid composition could alter nutritional and commercial values of milk as an input for dairy processing. Thus, milk fatty composition is of great importance to the dairy industry as most milk is traded as a processed product.

Keywords: Milk, milk fat, non-genetic factor, PUFA, human health

1. Introduction

Recently, the fatty acid composition in cattle milk has received considerable importance due to its major implications for human health and milk quality characteristics. Milk fat is one of the most complex natural fats that consist of approximately 400-500 fatty acids [1]. Short and medium chain fatty acids are synthesized *de novo* in the mammary gland, whereas, long chain fatty acids are derived from blood lipids, which are originating from the diet and endogenously produced lipids [2, 3]. Milk fat, either obtained from the circulating fat or synthesised in the mammary gland, is secreted as a mixture of milk fat globules (MFG) with different sizes, with a mean diameter between 3.3 and 5.2 μm , but ranging from 0.1 to 20 μm [4, 5]. Milk fatty acid composition traits are the most economically important traits controlled by numerous genes and environmental factors in dairy animals.

2. Fatty acid composition

Milk fat is characterized by high amounts of saturated fatty acids (SFA) and low amounts of polyunsaturated fatty acids (PUFA). A more unsaturated milk fat is preferred from human nutritional and health perspectives. Long chain fatty acid especially conjugated linoleic acid (CLA) have a role in modulation of plasma lipid concentration as well as anti-carcinogenic and anti-inflammatory effects [6]. Triglycerides, the lipidic fraction of milk contain ~70% saturated fatty acids (SFA) and 30% of unsaturated fatty acids (UFA) [7]. Low proportions of UFA and relatively high proportion of SFA have been shown to increase the level of blood LDL-cholesterol [8], especially C14:0 and C16:0, which is associated with both the increase of the blood cholesterol levels and cardiovascular disease in humans [9, 10].

In cattle milk PUFA account for as little as ~3% of all fatty acids [11]. The predominant n-3 FA in milk fat of the majority of mammals is α -linolenic acid. Among the MUFA group, the oleic acid (C18:1) is characterised by the highest content (24%), [12, 13]. O'Donnell *et al.* [14] and Strzałkowska *et al.* [15] reported that in ruminant's milk, there are also relatively small but significant contributions from other MUFA such as C14:1 (about 1%), C16:1 (about 1.5%), and very desirable vaccenic acid, which is a precursor of CLA in human organism (1.5-5%).

3. Factor affecting the milk composition

Milk fatty acid composition is variable in dairy cows and is influenced by species differences, parity or lactation order, age and size of the cow, dietary composition, feeding, season, pregnancy, sire service period, lameness, heat stress, health status of the cow, milking interval, genetic factors and other day to day variation behaviour or milking temperament, udder health, locality and stage of lactation. The effect of different non genetic factors affecting milk fatty acid composition are:

3.1. Effect of Breed /genetic group

Between and within breeds, fat varies the most and lactose the least [16]. Gaunt [17] reported cattle in the United States tend to have the lowest percentage of milk fat. The lower milk fat% of 4.73 ± 0.39 is seen in Gir cows [18]. The differences in the gene frequencies controlling the quality and quantity of milk components largely account for the average genetic differences among the breed. It has been found that the difference in milk constituents among the individuals within a breed is greater than the average difference among breed. The variation of fat percentage within a breed may be as much as 2%. Some Holstein Friesian (H.F.) produced milk contains 5% fat, whereas, some Jersey produced as low as H.F. breed average [19].

Poulsen *et al.* [20] showed that there was a considerable difference between breeds in the milk fatty acid composition. Some of the FA are more regulated by the environment (long chain FAs), while, the regulation of other FAs (short and medium chain FAs) is more influenced by genetics. The Danish Jersey (DJ) cows were characterized by higher levels of saturated short chain FAs compared to the Danish Holstein (DH), whereas, the DH had higher content of unsaturated C18 FA.

Lowest fat% (3.33 ± 0.03), in milk was reported by the Misra and Joshi [21] from Karan Fries cows during the rainy season in Karnal and highest fat% (5.98 ± 0.11) was reported by Yadav *et al.* [22] from JH cross bred in Hisar. Frequency distributions demonstrated that milk fat content of CLA and CLA-desaturase index varied over three fold among individuals. Holsteins had a higher milk fat content of CLA and CLA-desaturase index, but breed differences were minor [23]. Misra [24] reported that Jersey half breeds were superior to the H.F. half breeds with respect to fat% and SNF% in milk.

3.2. Effect of Season

The effects of seasonal variation on milk fatty acid composition and the physico-chemical properties vary throughout the year to different extents. Milk composition traits were highest in hot humid season but lowest in milk yield as compared to other seasons according to Sarkar *et al.* [25].

Lowest fat% (3.33 ± 0.03), in milk was reported by the Misra and Joshi [21] from Karan Fries cows during the rainy season in Karnal and highest fat% (5.98 ± 0.11) was reported by Yadav *et al.* [22] from JH cross bred in Hisar. DairyCo [26] reported that in the years 2009–2013, fat levels from the UK national herd gradually decreased from January to July, followed by a sharp increase to more than 4.20% in August and September, and remained constant in October, November and December.

Heck *et al.* [27] reported lower fat and protein contents in summer than in winter milk. This could be attributed to the different temperatures and feed composition, because cows consume more dry feed in winter, whereas, in summer they

eat grass and stay outside for longer [28].

Raw milk produced in the autumn period had a significantly higher fat content than in other periods which is broadly in line with the UK national statistics [26]. Yadav *et al.* [29] reported that fat percentage higher in winter calvers during 1st and 10th month of lactation, in the rainy season calver during 3rd and 4th lactation, in summer calvers 6th, 7th, and 8th month of lactation. Wiking *et al.* [30] showed that MFG sizes vary with diurnal fat production. As daily milk fat production increases, so does the milk fat globule size.

3.3. Effect of Parity/ Lactation

The age of the lactating animal is an important physiological factor affecting the composition of milk [31]. Barbary *et al.* [32] observed that milk fat percentage of H.F. cows was significantly affected by parity. There is report that milk fat% (4.9 ± 0.02) and SNF% (9.1 ± 0.01) contents of Sahiwal cows attained a peak value in the 3rd lactations and declined thereafter to the minimum ($4.8\pm 0.06\%$ fat and $9.0\pm 0.05\%$ SNF) in the 10th [33]. Misra and Joshi [21], furthermore, reported that the milk constituents had highest value in 1st parity, and whereas, the observation of Yadav *et al.* [22] revealed a gradual increase in the average fat% in milk during the whole lactation, on or up to peak day and on or up to week parity.

Lactation number of the animal has no significant effect on fat and SNF content of Tharparkar, Red Sindhi and Karan Swiss cow [33]. Shavaby [34] also found that the effect of lactation were non-significant for fat percentage in milk.

Schutz *et al.* [35] reported that fat yield was increased with parity of Guernsey, Holstein and Jersey cows. They also reported that sample day milk constituent's yields (including fat yield) were greater for Holsteins than Guernsey or Jerseys. The effect of parity was significant on fat% in Karan Fries cattle [21]. However, perusal of literature revealed both significant [36] and non-significant [37] effect of parity on different milk constituents and yield traits.

3.4. Effect of Stage of lactation

Generally, higher milk yield at early lactation are accompanied by low constituents percentage [38, 39]. Fat percentage was higher in 1st month of lactation. It decreased thereafter until the fifth or sixth month and increased subsequently in H.F. cows [40]. Altenhofer *et al.* [41] found significant difference for milk fat globules from the start, middle and end of lactation in bovine. Similar results have been observed in ewes [42].

Zurkowska [43] reported that milk fat% declined in the 1st month of lactation and in 2nd month it was maintained at a low level. Then at 3rd month it get increased. Intriери and Minieri [44] revealed that fat% decreased in the first two months and then gradually increased reaching to maximum towards the end of lactation. Sharma *et al.* [45] found that fat percentage declined initially, then increasing after 4 months in H.F. whereas, in Jerseys the rise was from the beginning of lactation.

4. Conclusion

The composition of cows' milk fat is influenced by numerous non genetic factors. Therefore, desirable changes in fat composition can be achieved by adjusting non genetic factors to improve the nutritional value of milk. The utilization of these factors could be possible owing to the genetic variability in fatty acid composition. It is important to understand the

relationship of non-genetic factor effects with milk production, resulting from genetic correlations between milk yield, fat content and proportions of FAs. Selection of cows for low fat content can result in a more desirable milk fat composition for human health, while selection for milk yield can affect the proportion of most individual FAs to a limited extent.

5. References

1. Barłowska J, Litwinczuk Z. Nutritional and pro-health properties of milk fat. *Medycyna Weterynaryjna*. 2009; 65:171-174.
2. Berquin IM, Edwards IJ, Chen YQ. Multi-targeted therapy of cancer by omega-3 fatty acids. *Cancer letters*. 2008; 269:363-377.
3. Brenna JT, Salem NJr, Sinclair AJ, Cunnane SC. α -Linolenic acid supplementation and conversion to n-3 long-chain polyunsaturated fatty acids in humans. *Prostaglandins, Leukotrienes and Essential Fatty Acids*. 2009; 80:85-91.
4. Menard O, Ahmad S, Rousseau F, Briard-Bion V, Gaucheron F, Lopez C. Buffalo vs. cow milk fat globules: size distribution, zeta-potential, compositions in total fatty acids and in polar lipids from the milk fat globule membrane. *Food Chemistry*. 2010; 120:544-551.
5. Lopez C. Milk fat globules enveloped by their biological membrane: Unique colloidal assemblies with a specific composition and structure. *Current Opinion in Colloid and Interface Science*. 2011; 16:391-404.
6. Haug A, Hostmark A, Harstad O. Bovine milk in human nutrition. *Lipids in Health and Disease*. 2007; 6:25.
7. Altenhofer C, Spornraft M, Kienberger H, Rychlik M, Herrmann J, Meyer HH *et al*. Effects of rapeseed and soybean oil dietary supplementation on bovine fat metabolism, fatty acid composition and cholesterol levels in milk. *Journal of Dairy Research*. 2014; 81:120-128.
8. Temme EHM, Mensink RP, Hornstra G. Comparison of the effects of diets enriched in lauric, palmitic or oleic acids on serum lipids and lipoproteins in healthy men and women. *The American Journal of Clinical Nutrition*. 1996; 63:897-903.
9. Mozaffarian D. Does alpha-linolenic acid intake reduce the risk of coronary heart disease? A review of the evidence. *Alternative Therapies in Health and Medicine*. 2005; 11:24-30.
10. Stoop WM, van Arendonk JA, Heck JM, van Valenberg HJ, Bovenhuis H. Genetic parameters for major milk fatty acids and milk production traits of Dutch Holstein-Friesians. *Journal of Dairy Science*. 2008; 91(1):385-394.
11. Devle H, Rukke EO, Naess-Andresen CF, Ekeberg D. A GC-magnetic sector MS method for identification and quantification of fatty acids in ewe milk by different acquisition modes. *Journal of Separation Science*. 2009; 32:3738-3745.
12. Schmidely P, Andrade PVD. Dairy performance and milk fatty acid composition of dairy goats fed high or low concentrate diet in combination with soybeans or canola seed supplementation. *Small Ruminant Research*. 2011; 99:135-142.
13. Szumacher-Strabel M, Cieslak A, Zmora P, Pers-Kamczyc E, Bielińska S, Stanisław M *et al*. *Camelina sativa* cake improved unsaturated fatty acids in ewe's milk. *Journal of the Science of Food and Agriculture*. 2011; 91:2031-2037.
14. O'Donnell AM, Spatny KP, Vicini JL, Bauman DE. Survey of the fatty acid composition of retail milk differing in label claims based on production management practices. *Journal of Dairy Science*. 2010; 93:1918-1925.
15. Strzałkowska N, Jozwik A, Bagnicka E, Krzyżewski J, Horbanczuk JO. Studies upon genetic and environmental factors affecting the cholesterol content of cow milk. II. Effect of silage type offered. *Animal Science Papers and Reports*. 2009; 27:199-206.
16. Woodford JA, Jorgensen NA, Barrington GP. Impact of dietary fiber and physical form on performance of lactating dairy cows. *Journal of Dairy Science*. 1986; 69:1035-1047.
17. Gaunt SN. Genetic variation in the yields and contents of milk constituents. *Bulletin of the International Dairy Federation*. 1980; 125:73-82.
18. Padekar RN, Bhoite UY. Effect of stage of lactation on milk yield and its constituents in Gir cows. *Indian Veterinary Journal*. 2002; 79:181-182.
19. Webb BH, Johnson AH, Alfred JA. Fundamentals of dairy chemistry. 2nd edition. The AVI publication company. INV Western Port Connecticut, 1974.
20. Poulsen NA, Gustavsson F, Glantz M, Paulsson M, Larsen LB, Larsen MK. The influence of feed and herd on fatty acid composition in 3 dairy breeds (Danish Holstein, Danish Jersey, and Swedish Red). *Journal of Dairy Science*. 2012; 95:6362-6371.
21. Mishra SS, Joshi BK. Genetic and non-genetic factors affecting lactation milk constituents and yield traits in Karan Fries cattle. *Indian Journal of Dairy Sciences*. 2004; 57(1):69-72.
22. Yadav SBS, Yadav AS, Yadav BI, Yadav MS. Factors affecting fat percentage in crossbred dairy cattle. *Indian Journal of dairy science*. 1989; 42(3):475-481.
23. Kelsey JA, Corl BA, Collier RJ, Bauman DE. The effect of breed, parity, and stage of lactation on conjugated linoleic acid (CLA) in milk fat from dairy cows. *Journal of Dairy Science*. 2003; 86(8):2588-2597.
24. Misra RK. Milk production and milk composition of Jersey half-bred and Holstein half-bred cows. *Indian Journal of Animal health*. 2000; 39(1):1-4.
25. Sarkar U, Gupta AK, Sarkar V, Mohanty TK, Raina VS, Prasad S. Factors affecting test day milk yield and milk composition in dairy animals. *Journal of dairying, foods & home sciences*. 2006; 25(2):129-132.
26. Dairy Co. MILKBENCH Report. Agriculture and Horticulture Development Board Stoneleigh Park Kenilworth, Warwickshire CV8 2TL, UK, 2013.
27. Heck JML, Van Valenberg HJF, Dijkstra J, Van Hooijdonk ACM. Seasonal variation in the Dutch bovine raw milk composition. *Journal of Dairy Science*. 2009; 92:4745-4755.
28. Fox PF, McSweeney, Paul LH. *Advanced Dairy Chemistry: Volume 1: Proteins, Parts A & B*. 3rd edition. 2003, 480-481.
29. Yadav PL, Nautiyal LP, Saxena MM, Sanyal MK, Dubey PC, Pandey HS *et al*. Quality characteristics of milk from Holstein Friesian and Hariana x exotic cattle. *Indian veterinary Journal*. 1991; 68:963-966.
30. Wiking L, Stagsted J, Bjorck L, Nielsen JH. Milk fat globule size is affected by fat production in dairy cows. *International Dairy Journal*. 2004; 14:909-913.
31. Singh A, Gandhi RS, Chakarvarty AK, Sharma RC,

- Gurnani M. Effect of postpartum breeding interval on milk production and reproduction traits in Karan Fries cattle. *Indian Journal of Animal Research*. 1979; 35(2):83-87.
32. Barbary EIASA, Mahdy AE, Shafie OM, Aly HM. Some factor affecting milk production and milk constituents and their relation to udder measurements in pure Friesian cows. *Alexandria Journal of Agricultural Research*. 1999; 44(1):17-35.
33. Lal D, Narayan KM. Effect of lactation number of the animal on the fat and solid-not-fat contents of milk. *Indian Journal of Animal Science*. 1984; 54(9):835-839.
34. Shavaby MA. Factors influencing the concentrations and yields of milk constituents and their inter-relationship. *Journal of Dairy Research*. 1988; 55(2):171-177.
35. Schutz MM, Hansen LB, Steuernagel GR, Kuck AL. Variation of milk, fat, protein and somatic cells for Dairy cattle. *Journal of Dairy Science*. 1990; 73:484-494.
36. Jain DK, Sharma KNS, Bhatnagar DS, Sharma RC. Fat and SNF concentration in cow milk as affected by parity and stage of lactation. *Indian Journal of Dairy Science*. 1987; 40(1):111-118.
37. Saxena MM, Suman CL, Katpatal BG, Pandey HS. Study of milk constituents and their yield in Holstein-Friesian cows. *Indian Journal of Animal Production and Management*. 1997; 13(3):127-130.
38. Eftala E. Factors affecting the composition of milk. II. Effect of stage and number of lactation and some external factors and significance of choice of test days in estimation of protein producing ability. *Annales Agriculturae Fenniae*. 1976; 15:196.
39. Freeman TR. Effect of breed, season, and stage of lactation on certain constituents and properties of milk. *Kentucky Agricultural Experiment Station. University Kentucky, Lexington*. 1959, 667.
40. Tamarcenko ME, Finkelj SB, Cibizov NI, Lebedev DS. The determination of fat-content in cow's milk for a lactation. *Animal Breeding Abstract*. 1950; 19:601.
41. Altenhofer C, Holzmuller W, Wolfertstetter F, Ribeiro DW, Kulozik U, Pfaffl MW *et al*. Temporal variation of milk fat globule diameter, fat and cholesterol content and milk epithelial cell gene expression in dairy cows. *International Journal of Dairy Technology*. 2015; 68:519-526.
42. Martini M, Altomontel I, Pesi R, Tozzi MG, Salari F. Fat globule membranes in ewe's milk: the main enzyme activities during lactation. *International dairy journal*. 2012; 28(1):36-39.
43. Zurkowska K. Investigations on the content of protein and fat in the milk of Black piedlow land cattle. *Animal Breeding Abstract*. 1966; 35:3432.
44. Intriери F, Minieri L. The chemical composition of milk of buffalos in Caserta province in relation to interval from calving. *Animal Breeding Abstract*. 1967; 36:2377.
45. Sharma AK, Wilcox CJ, Martin FG, Thatcher WW. Effect of stage of lactation and pregnancy and their interactions on milk yield and constituents. *Journal of Dairy Science*. 1990; 73:1586-1592.