



ISSN 2320-7078
JEZS 2014; 2 (3): 164-169
© 2014 JEZS
Received: 29-05-2014
Accepted: 30-06-2014

Xin Chao Liu
Department of Entomology, China
Agricultural University, Haidian
District, Beijing 100193, China.

Jin Feng Hu
Institute of Plant Protection, Fujian
Academy of Agricultural Sciences,
Fuzhou, Fujian 350013, China

Ligang Zhou
Department of Plant Pathology, China
Agricultural University, Haidian
District, Beijing 100193, China

Zhi Long Liu
Department of Entomology, China
Agricultural University, Haidian
District, Beijing 100193, China

Correspondence:
Zhi Long Liu
Department of Entomology, China
Agricultural University, Haidian
District, Beijing 100193, China

Evaluation of fumigant toxicity of essential oils of Chinese medicinal herbs against *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae)

Xin Chao Liu, Jin Feng Hu, Ligang Zhou and Zhi Long Liu

ABSTRACT

Essential oils of sixteen Chinese medicinal plants were screened for fumigant toxicity against *Bemisia tabaci*. The essential oil of *Allium sativum* exhibited the strongest fumigant activity against *B. tabaci* with an LC₅₀ value of 0.11 µg/l followed by the essential oils of *Agastache rugosa* (LC₅₀ = 7.08 µg/l), *Illicium verum* (LC₅₀ = 8.61 µg/l), *Chenopodium ambrosioides* (LC₅₀ = 8.80 µg/l), and *Syzygium aromaticum* (LC₅₀ = 13.54 µg/l). Three essential oils (*Curcuma aeruginosa*, *Schizonepeta tenuifolia* and *Valeriana officinalis*) did not show fumigant toxicity at a concentration of 50.00 µg/l. Chemical composition of garlic essential oil was determined using GC-MS. Two main constituents of the garlic oil, diallyl trisulfide and diallyl disulfide exhibited strong fumigant toxicity against the whitefly with LC₅₀ values of 0.08 µg/l and 0.12 µg/l, respectively while diallyl sulfide has a LC₅₀ value of 1.68 µg/l.

Keywords: *Bemisia tabaci*; *Allium sativum*; fumigation; essential oil composition, allyl trisulfide; diallyl disulfide.

1. Introduction

The sweet potato whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae) is a major pest of economically important crops worldwide [1]. In China, it has been recorded since 1940's, but was not considered as a serious pest until recent years [2]. However, since 1996 it has been regarded as a serious pest of cotton, vegetables and ornamental crops in 22 provinces or cities in mainland China. Vegetable growers observed regular outbreaks and up to 70% losses in cucumber from *B. tabaci* infections in the Beijing districts [2]. The development of resistance of this pest to existing conventional pesticides and the increasing public concern over environmental pollution and health hazards created by synthetic pesticides generate a great need for new types of pest control agent's advantage with higher activity against the target pests, and lower impact on humans and environmental quality. The use of biologically based compounds in plant extracts or essential oils may be an alternative to currently used insecticides to control insects [3,4]. Moreover, essential oils have a broad spectrum of insecticidal activity due to the presence of several modes of action, including repellent and antifeedant activities, inhibition of molting and respiration, reduction in growth and fecundity, cuticle disruption and effect on the invertebrate octopamine pathway [3,5]. Essential oils derived from plants may also have minimal direct and/or indirect effects on natural enemies [6,7]. Essential oils and aqueous extracts of some medicinal plants have been evaluated for insecticidal activity against the sweet potato whitefly (*B. tabaci*) and some of them show potential to be developed as natural insecticides in the control of the whitefly [8-22]. Several essential oils derived from Chinese medicinal herbs have been screened for repellency and insecticidal activities against grain storage insects and cockroaches as well as mosquitoes [23-28]. However, little information on fumigant activity of the essential oils of Chinese medicinal herbs against the sweet potato whitefly was available. In this study, we assessed fumigant activity of 16 essential oils derived from Chinese medicinal herbs/spices against the whitefly, *B. tabaci*. Moreover, the two main constituent compounds of garlic essential oil (the strongest fumigant toxicity among the essential oils tested) were also evaluated for fumigant activity against the sweet potato whitefly.

2. Materials and Methods

2.1 Insects

Colonies of the sweet potato whitefly, *B. tabaci* were established from individuals collected from sweet potato plants in a greenhouse (25-28 °C) of the Institute of Plant Protection, Fujian

Academy of Agricultural Sciences, Fuzhou, China and were maintained on sweet potato plants. To obtain uniform age in insects for bioassay studies, vigorous *B. tabaci* adults were selected from the population to produce eggs on potted, caged, insect-free plants for 36 h. The adults were removed to allow eggs of uniform age to develop in a growth chamber at 24.5-25.5 °C, 70-75 relative humidity (R.H.) and 16:8 h light: dark photoperiod for a certain period to obtain vigorous two-day-old adults.

2.2. Chinese medicinal herbs

Garlic (*Allium sativum*) was purchased fresh from a local supermarket in Beijing. The other medicinal herbs (Table 1) were obtained from Guangming Chinese Medicinal Herb Decoction Company (Hebei 071200, China). All the medicinal herbs were identified by Dr. Liu, QR (College of Life Sciences, Beijing Normal University, Beijing, China) and the voucher specimens were deposited in the museum of Department of Entomology, China Agricultural University, Beijing, China.

Table 1: Chinese medicinal herbs used for fumigant activity

Botanical name	Common name	Family	Plant material
<i>Acorus tatarinowii</i> Schott	Sweet flag	Araceae	Rhizomes
<i>Agastache rugosa</i> (Fisch et Mey) O. Ktze	Korean Mint	Labiatae	Aerial parts
<i>Allium sativum</i> L.	Garlic	Lilliaceae	Bulbs
<i>Alpinia officinarum</i> Hance	Lesser galangal	Zingiberaceae	Rhizomes
<i>Cuminum cyminum</i> L.	Cumin	Umbelliferae	Fruits
<i>Curcuma aeruginosa</i> Roxb.	Pink and blue ginger	Zingiberaceae	Rhizomes
<i>Chenopodium ambrosioides</i> L.	Mexican tea	Chenopodiaceae	Aerial parts
<i>Cnidium monnieri</i> (L.) Cusson	Monnier's snow parsley	Umbelliferae	Seeds
<i>Elsholtzia ciliata</i> (Thunb.) Hyland	Vietnamese balm	Labiatae	Aerial parts
<i>Foeniculum vulgare</i> Mill.	fennel	Umbelliferae	Fruits
<i>Illicium verum</i> Hook.f.	Star anise	Magnoliaceae	Fruits
<i>Schizonepeta tenuifolia</i> (Benth.) Briq.	Japanese mint	Labiatae	Flowers
<i>Syzygium aromaticum</i> (L.) Merr. et Perry	Cloves	Myrtaceae	Flower buds
<i>Valeriana officinalis</i> L.	Valerian	Valerianaceae	Rhizomes
<i>Zanthoxylum bungeanum</i> Maxim.	Prickly ash	Rutaceae	Fruits
<i>Z. schinifolium</i> Sieb. et Zucc	Green prickly ash	Rutaceae	Fruits

2.3. Essential oil extraction

Fresh garlic and lesser galangal and other dried spices (grinded to powdered form using a grinding mill) were subjected to hydrodistillation using a modified Clevenger-type apparatus for 6 h and extracted with *n*-hexane. Anhydrous sodium sulfate was used to remove water after extraction. Essential oils were stored in airtight containers in a refrigerator at 4 °C for subsequent experiments. Diallyl disulfide, diallyl trisulfide and diallyl sulfide were purchased from Aladdin-reagent Company (Shanghai, China).

2.4. Fumigant toxicity

Desiccators with a capacity of 4l were used as test chambers. Adults of *B. tabaci* on fresh leaves of sweet potato were exposed separately to the essential oils of Chinese medicinal herbs and constituent compounds of garlic oil, respectively. The leaves infested with *B. tabaci* were brought to desiccators. In order to maintain turgor of leaves, their petioles were dipped through a hole made in a

rubber cap fitted to 15 ml capacity glass vials filled with tap water. The control consisted of a similar setup but without essential oils/compounds. Each replicate consisted of 20 *B. tabaci* adults placed on one leaf. For each dose, five replicates were used. The essential oils/compounds were applied with an automatic pipette on a blotting paper strip (6 cm×3 cm) attached to the bottom of the desiccators. Initial testing was establishing appropriate dose ranges and the six concentrations were used. No material was applied to the control desiccators. Mortality of insects was observed after 24 h.

2.5. GC-MS analysis

Gas chromatographic analysis was performed on an Agilent 6890N Gas Chromatograph (Agilent Technologies, Santa Clara, USA) while the essential oils were identified on an Agilent Technologies 5973N mass spectrometer. It was equipped with a flame ionization detector and HP-5MS (Agilent Technologies, Santa Clara, USA) capillary column (30 m × 0.25 mm × 0.25 μm). The GC settings were as

follows: the initial oven temperature was held at 60 °C for 1 min and increased at 10 °C/min to 180 °C, hold for 1 min, and then ramped at 20 °C/min to 280 °C, hold for 15 min. The injector temperature was maintained at 270 °C. The samples (1 µl, diluted 1:100 in acetone) were injected, with a split ratio of 1:10. The carrier gas was helium at a flow rate of 1.0 ml/min. Spectra were scanned from m/z 20 to 550 at 2 scans/s. Most constituents were identified using gas chromatography by comparison of their retention indices with those of the literature or with those of authentic compounds available in our laboratories. The retention indices were determined in relation to a homologous series of *n*-alkanes (C₈–C₂₄) under the same operating conditions. Further identification was made by comparison of their mass spectra on both columns with those stored in NIST 05 and Wiley 275 libraries or with mass spectra from literature [29]. Component relative percentages were calculated on GC peak areas.

2.6. Statistical analysis

The observed mortality data were corrected for control mortality using Abbott's formula. The results from all replicates were subjected to Probit analysis using PriProbit Program V1.6.3 to determine LC₅₀ values with their fiducial limits [30]. Samples for which the 95 % fiducial limits did not overlap were considered to be significantly different.

3. Results and Discussion

3.1. Chemical composition of the essential oil

The yield of *A. sativum* essential oil was 1.83% (V/W) and the density of the essential oil was determined to be 1.05 g/ml. The chemical composition of the essential oil was summarized in Table 2. A total of 17 components of the essential oil were identified, accounting for 98.90% of the total oil. The principal compound in the essential oil was diallyl trisulfide (61.87%) followed by diallyl disulfide (19.47%), diallyl sulfide (4.41%) and diallyl tetrasulfide (3.67%).

Table 2: Chemical constituents of the essential oil derived from *Allium sativum* bulbs

Peak no.	Compounds	RI	Composition (%)
1	Dimethyl disulfide*	740	0.28
2	Diallyl sulfide*	848	4.41
3	Allyl isothiocyanate	890	0.49
4	Allyl methyl disulfide	915	1.46
5	Dimethyl trisulfide	975	1.38
6	1,3-Dithiane	1027	0.84
7	Diallyl disulfide*	1077	19.47
8	Linalool*	1097	0.11
9	Allyl methyl trisulfide	1134	0.69
10	Borneol*	1165	0.37
11	Methyl propyl trisulfide	1168	0.53
12	α-Terpineol*	1188	1.02
13	Dimethyl tetrasulfide	1224	0.24
14	Diallyl trisulfide*	1296	61.87
15	Diallyl thiosulfinate	1325	1.04
16	Allyl methyl tetrasulfide	1386	1.03
17	Diallyl tetrasulfide	1540	3.67
	Total		98.90

RI, retention index as determined on a HP-5MS column using the homologous series of *n*-hydrocarbons. *Identification by co-injection of authentic compounds.

3.2. Fumigant toxicity of the essential oils

Among 16 essential oils tested (Table 3), the essential oil of *A. sativum* exhibited the strongest fumigant activity against *B. tabaci* adults with an LC₅₀ value of 0.11 µg/l followed by the essential oils of *A. rugosa* (LC₅₀ = 7.08 µg/l), *I. verum* (LC₅₀ = 8.61 µg/l), *C. ambrosioides* (LC₅₀ = 8.80 µg/l), *S. aromaticum* (LC₅₀ = 13.54 µg/l) and *C. monnieri* (LC₅₀ = 19.31 µg/l) (Table 3). The other essential oils of *A. tatarinowii*, *Z. bungeanum*, *Z. schinifolium*, *A. officinarum*, *E. ciliate*, and *C. cyminum* also possessed fumigant toxicity against *B. tabaci* adults with LC₅₀ values ranged from 21.83 µg/l to 31.17 µg/l. However, the other three essential oils (*C. aeruginosa*, *S. tenuifolia* and *V. officinalis*) did not show fumigant toxicity at a concentration of 50.00 µg/l. The essential oil of *A. sativum* possessed almost 64 times

stronger fumigant activity than the second active essential oil of *A. rugosa* (Table 3). In previous reports, the essential oil of *A. sativum* possessed fumigant activity against adult German cockroaches, *Blattella germanica* [31], larval sciarid fly, *Lycoriella ingenua* [32], the Japanese termite (*Reticulitermes speratus*) [33], pear pear psylla (*Cacopsylla chinensis*) [34], and several grain storage insects (*Ephestia kuehniella*, *Sitophilus oryzae*, *S. zeamais* and *Tribolium castaneum*) [35-37]. The essential oils of *A. rugosa*, *I. verum*, *C. ambrosioides*, and *S. aromaticum* have been also demonstrated to possess fumigant activity against several insect pests, e.g. head lice (*Pediculus humanus capitis*) [38], larval sciarid fly, *L. ingenua* [32], adult cigarette beetle (*Lasioderma serricornis*) [39], adults and larvae of the pulse beetle (*Callosobruchus chinensis*) [40], and adults of the

maize weevil (*S. zeamais*)^[41].

Table 3: Fumigant toxicity of the essential oils derived from the Chinese medicinal herbs and constituent compounds of garlic oil against the adult whitefly

Treatment	LC ₅₀ (µg/l air)	95% fiducial limits	Slope ± SD	Chi square (χ ²)
<i>A. tatarinowii</i>	21.83	19.95-24.07	1.33 ± 0.18	17.48
<i>A. rugosa</i>	7.08	5.88-8.24	1.85 ± 0.11	18.61
<i>A. sativum</i>	0.11	0.09-0.14	1.23 ± 0.11	7.89
<i>A. officinarum</i>	25.77	22.89-28.27	2.78 ± 0.17	13.08
<i>C. cyminum</i>	30.17	26.98-33.98	9.58 ± 0.85	13.12
<i>C. aeruginosa</i>	>50.00	-	-	-
<i>C. ambrosioides</i>	8.85	8.16-9.87	2.70 ± 0.15	18.65
<i>C. monnieri</i>	19.31	18.04-21.26	2.13 ± 0.12	9.45
<i>E. ciliata</i>	26.64	24.23-28.54	2.56 ± 0.19	12.57
<i>F. vulgare</i>	10.86	9.50-12.35	2.87 ± 0.21	18.34
<i>I. verum</i>	8.61	7.34-9.41	2.68 ± 0.19	13.89
<i>S. tenuifolia</i>	>50.00	-	-	-
<i>S. aromaticum</i>	13.54	11.93-14.79	2.70 ± 0.19	13.57
<i>V. officinalis</i>	>50.00	-	-	-
<i>Z. bungeanum</i>	23.76	21.55-26.16	4.78 ± 0.27	12.36
<i>Z. schinifolium</i>	24.37	20.65-26.55	4.12 ± 0.29	9.12
diallyl disulfide	0.12	0.11-0.14	1.37 ± 0.09	9.37
diallyl trisulfide	0.08	0.07-0.09	1.19 ± 0.10	12.34
diallyl sulfide	1.68	1.55-1.81	1.78 ± 0.19	13.45

LC₅₀, median lethal concentration; SD, standard deviation

The above findings suggest that the fumigant toxicities of these mentioned essential oils of Chinese medicinal herbs especially the essential oil of *A. sativum* are quite promising. They showed potential to be developed as possible natural fumigants for the control of the whitefly by considering the currently used synthetic fumigants (insecticides) because of being highly toxic to humans and other non-target organisms. Moreover, most of these Chinese medicinal herbs are widely cultivated and easily grown. However, for the practical application of the essential oils and their constituent compounds as novel fumigants, further studies on the safety of the essential oils/compounds to humans and plants are needed. A further study is also necessary to determine the toxicity of these essential oils and their constituent compounds on other economically important pests and their natural enemies in greenhouse conditions where pest management depends on chemical applications. Further studies on the development of formulations are also necessary to improve the efficacy and stability and to reduce cost.

3.3. Fumigant toxicity of the constituents

The two main constituents of *A. sativum* oil, diallyl trisulfide and diallyl disulfide exhibited strong fumigant toxicity against the whitefly *B. tabaci* adults with LC₅₀ values of 0.08 µg/l and 0.12 µg/l, respectively, while diallyl sulfide has a LC₅₀ value of 1.68 µg/l. Diallyl trisulfide shows stronger fumigant activity than the crude essential oil and diallyl disulfide. Moreover, diallyl trisulfide exhibited 20 times stronger fumigant toxicity against the whitefly than another constituent compound, diallyl sulfide (Table 3). It suggested that the fumigant activity of the essential oil of *A. sativum* may be attributed to diallyl trisulfide. In the

previous reports, diallyl trisulfide has been demonstrated to exhibit strong fumigant activity against the maize weevil (*S. zeamais*), rice weevil (*S. oryzae*) and red flour beetle (*T. castaneum*)^[35, 42]. Diallyl trisulfide was also shown to have stronger fumigant toxicity than diallyl disulfide against the Japanese termite (*R. speratus*)^[31]. Huang *et al.*^[35] also demonstrated that diallyl trisulfide was strongest fumigant constituent in the essential oil of *A. sativum*. However, Tunaz *et al.*^[31] found that allyl isothiocyanate derived from the essential oil of *A. sativum* exhibited fumigant toxicity against adult German cockroaches (*B. germanica*) with an LC₅₀ value of 0.68 µl/l and allyl isothiocyanate also exhibited stronger fumigant activity than diallyl disulfide against larval sciarid fly (*L. ingenua*)^[32]. However, allyl isothiocyanate was not detected from the essential oil of *A. sativum* in present study. Allyl isothiocyanate mainly comes from the cruciferous vegetables, e.g. seeds of black mustard (*Brassica nigra*) or brown Indian mustard (*Brassica juncea*) and only minor components in the essential oil of *A. sativum*^[35, 43]. Moreover, diallyl disulfide derived from garlic essential oil exhibited strong larvicidal activity against *Culex pipiens* larvae^[44].

4. Conclusions

Thirteen of sixteen essential oils of Chinese medicinal herbs and three main constituents of the essential oil of *A. sativum* demonstrated some fumigant toxicity against the whitefly. They showed potential to be developed as possible natural fumigants for control of the whitefly but needs to be further evaluated for safety in humans and to enhance its activity.

5. Acknowledgments

This project was supported by the Hi-Tech Research and

Development of China 2011AA10A202. We thank Liu Q.R. from the College of Life Sciences, Beijing Normal University, Beijing 100875, for the identification of the investigated medicinal herbs

6. References

- Sertkaya E, Kaya K, Soylu S. Chemical compositions and insecticidal activities of the essential oils from several medicinal plants against the cotton whitefly, *Bemisia tabaci*. *Asian J Chem* 2010; 22:2982-2990.
- Wang LD, Huang J, You MS, Guan X, Liu B. Toxicity and feeding deterrence of crude toxin extracts of *Lecanicillium lecanii* (Hyphomycetes) against sweet potato whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae). *Pest Manag Sci* 2007; 63:381-387.
- Isman MB. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu Rev Entomol* 2006; 51:45-66.
- Regnault-Roger C, Vincent C, Arnason JT. Essential oils in insect control: Low-risk products in a high-stakes world. *Annu Rev Entomol* 2012; 57:405-424.
- Enan E. Insecticidal activity of essential oils: octopaminergic sites of action. *Comp Biochem Physiol* 2001; 130C:325-337.
- Bostanian NJ, Akalach M, Chiasson H. Effects of a Chenopodium-based botanical insecticide/acaricide on *Orius insidiosus* (Hemiptera: Anthocoridae) and *Aphidius colemani* (Hymenoptera: Braconidae). *Pest Manag Sci* 2005; 61:979-984.
- Cloyd RA, Galle CL, Keith SR, Kalscheur NA, Kemp KE. Effect of commercially available plant-derived essential oil products on arthropod pests. *J Econ Entomol* 2009; 102:1567-1579.
- Fenigstein A, Eliyahu M, Gan-Mor S, Veierov D. Effects of five vegetable oils on the sweetpotato whitefly *Bemisia tabaci*. *Phytoparasitica* 2001; 29:197-206.
- Kostyukovsky M, Ravid U, Shaaya E. The potential use of plant volatiles for the control of stored product insects and quarantine pests in cut flowers. *Acta Horticulturae* 2002; 576:347-358.
- Aslan I, Oezbek H, Calmasur O, Sahin F. Toxicity of essential oil vapors to two greenhouse pests, *Tetranychus urticae* Koch and *Bemisia tabaci* Genn. *Ind Crop Prod* 2004; 19:167-173.
- Calmasur O, Aslan I, Sahin F. Insecticidal and acaricidal effect of three Lamiaceae plant essential oils against *Tetranychus urticae* Koch and *Bemisia tabaci* Genn. *Ind Crop Prod* 2006; 23:140-146.
- Ateyyat MA, Al-Mazra'awi MT, Shatnawi MA. Aqueous extracts of some medicinal plants are as toxic as imidacloprid to the sweet potato whitefly, *Bemisia tabaci*. *J Insect Sci* 2009; 9:15.
- Sayeda FF, Torkey HM, Abou-Yousef HM. Natural extracts and their chemical constituents in relation to toxicity against whitefly (*Bemisia tabaci*) and aphid (*Aphis craccivora*). *Aust J Basic Appl Sci* 2009; 3:3217-3223.
- Lin CY, Wu DC, Yu JZ, Chen BH, Wang CL, Ko WH. Control of silverleaf whitefly, cotton aphid and kanzawa spider mite with oil and extracts from seeds of sugar apple. *Neotrop Entomol* 2009; 38:531-536.
- Ribeiro NC, Gomes CCA, Born FS, Abreu SHA. Insecticidal activity against *Bemisia tabaci* biotype B of peel essential oil of *Citrus sinensis* var. *pear* and *Citrus aurantium* cultivated in northeast Brazil. *Nat Prod Commun* 2010; 5:1819-1822.
- Yang NW, Li AL, Wan FH, Liu WX, Johnson D. Effects of plant essential oils on immature and adult sweetpotato whitefly, *Bemisia tabaci* biotype B. *Crop Prot* 2010; 29:1200-1207.
- Tia EV, Adima AA, Niamke SL, Jean GA, Martin T, Lozano P, Menut C. Chemical composition and insecticidal activity of essential oils of two aromatic plants from Ivory Coast against *Bemisia tabaci* G. (Homoptera: Aleyrodidae). *Nat Prod Commun* 2011; 6:1183-1188.
- Kim SI, Chae SH, Youn HS, Yeon SH, Ahn YJ. Contact and fumigant toxicity of plant essential oils and efficacy of spray formulations containing the oils against B- and Q-biotypes of *Bemisia tabaci*. *Pest Manag Sci* 2011; 67:1093-1099.
- Zandi-Sohani N. Efficiency of Labiateae plants essential oils against adults of cotton whitefly (*Bemisia tabaci*). *Indian J Agric Sci* 2011; 81:1164-1167
- Baldin ELL, Crotti AEM, Wakabayashi KAL, Silva JPGF, Aguiar GP, Souza ES, Veneziani RCS, Groppo M. Plant-derived essential oils affecting settlement and oviposition of *Bemisia tabaci* (Genn.) biotype B on tomato. *J Pest Sci* 2013; 86:301-308.
- Tia EV, Lozano P, Menut C, Lozano YF, Martin T, Niamke S, Adima AA. Potentiality of essential oils for control of the whitefly *Bemisia tabaci* Genn., a greenhouse pest. *Phytotherapie* 2013; 11:31-38.
- Chae SH, Kim SI, Yeon SH, Perumalsamy H, Ahn YJ. Fumigant toxicity of summer savory and lemon balm oil constituents and efficacy of spray formulations containing the oils to B- and neonicotinoid-resistant Q-biotypes of *Bemisia tabaci* (Homoptera: Aleyrodidae). *J Econ Entomol* 2014; 107:286-292.
- Liu ZL, Goh SH, Ho SH. Screening of Chinese medicinal herbs for bioactivity against *Sitophilus zeamais* Mostchulsky and *Tribolium castaneum* (Herbst). *J Stored Prod Res* 2007; 43:290-296.
- Liu ZL, Yu M, Li XM, Wan T, Chu SS. Repellent activity of eight essential oils of Chinese medicinal herbs to *Blattella germanica* L. *Rec Nat Prod* 2011; 5:176-183.
- Zhao NN, Zhou L, Liu ZL, Du SS, Deng ZW. Evaluation of toxicities of some common spices essential oils from China against *Liposcelis bostrychophila*. *Food Control* 2012; 26:486-490
- Liang Y, Li JL, Xu S, Zhao NN, Zhou L, Cheng J, Liu ZL. Repellent constituents of the essential oil of *Epimedium pubescens* aerial parts against two grain storage insects. *J Econ Entomol* 2013; 106:513-519.
- Liu XC, Dong HW, Zhou L, Du SS, Liu ZL. Essential oil composition and larvicidal activity of *Toddalia asiatica* roots against the mosquito *Aedes albopictus* (Diptera: Culicidae). *Parasitol Res* 2013; 112:1197-1203.
- Liu XC, Liu QY, Zhou L, Liu ZL. Evaluation of larvicidal activity of the essential oil of *Allium macrostemon* Bunge and its selected major constituent

- compounds against *Aedes albopictus* (Diptera: Culicidae). Parasite Vector 2014; 7:184.
29. Adams RP. *Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy*; Allured: Carol Stream, IL, USA, 2007.
 30. Sakuma M. Probit analysis of preference data. Appl Entomol Zool 1998; 33:339-347.
 31. Tunaz H, Er MK, Isikber AA. Fumigant toxicity of plant essential oils and selected monoterpenoid components against the adult German cockroach, *Blattella germanica* (L.) (Dictyoptera: Blattellidae). Turk J Agric Forest 2009; 33:211-217.
 32. Park IK, Choi KS, Kim DH, Choi IH, Kim LS, Bak WC, Choi JW, Shin SC. Fumigant activity of plant essential oils and components from horseradish (*Armoracia rusticana*), anise (*Pimpinella anisum*) and garlic (*Allium sativum*) oils against *Lycoriella ingenua* (Diptera: Sciaridae). Pest Manag Sci 2006; 62:723-728.
 33. Park IK, Shin SC. Fumigant activity of plant essential oils and components from garlic (*Allium sativum*) and clove bud (*Eugenia caryophyllata*) oils against the Japanese termite (*Reticulitermes speratus* Kolbe). J Agric Food Chem 2005; 53:4388-4392.
 34. Zhao NN, Zhang H, Zhang XC, Luan XB, Zhou C, Liu QZ, Shi WP, Liu ZL. Evaluation of toxicity of the essential oil of aged garlic (*Allium sativum*) bulbs and its constituent compounds against overwintering *Cacopsyllia chinensis* (Homoptera: Psyllidae). J Econ Entomol 2013; 106(3):1349-1354.
 35. Huang Y, Chen SX, Ho SH. Bioactivities of methyl allyl disulfide and diallyl trisulfide from essential oil of garlic to two species of stored-product pests, *Sitophilus zeamais* (Coleoptera: Curculionidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae). J Econ Entomol 2000; 93:537-543.
 36. Kim JH, Park IK. Fumigant toxicity of Korean medicinal plant essential oils and components from *Asiasarum sieboldi* root against *Sitophilus oryzae* L. Flav Fragr J 2008; 23:79-83.
 37. Mikhael AA. Potential of some volatile oils in protecting packages of irradiated wheat flour against *Ephesia kuehniella* and *Tribolium castaneum*. J Stored Prod Res 2011; 47:357-364.
 38. Toloza AC, Zygadlo J, Cueto GM, Biurrun F, Zerba E, Picollo MI. Fumigant and repellent properties of essential oils and component compounds against permethrin-resistant *Pediculus humanus capitis* (Anoplura: pediculidae) from Argentina. J Med Entomol 2006; 43:889-895.
 39. Kim SI, Park C, Ohh MH, Cho HC, Ahn YJ. Contact and fumigant activities of aromatic plant extracts and essential oils against *Lasioderma serricorne* (Coleoptera: Anobiidae). J Stored Prod Res 2003; 39:11-19.
 40. Chaubey MK. Fumigant toxicity of essential oils from some common spices against pulse beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). J Oleo Sci 2008; 57:171-179.
 41. Chu SS, Hu JF, Liu ZL. Composition of essential oil of Chinese *Chenopodium ambrosioides* and insecticidal activity against maize weevil, *Sitophilus zeamais*. Pest Manag Sci 2011; 67:714-718.
 42. Koul O. Biological activity of volatile di-n-propyl disulfide from seeds of neem, *Azadirachta indica* (Meliaceae), to two species of stored grain pests, *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst). J Econ Entomol 2004; 97:1142-1147.
 43. Wu H, Liu XR, Yu DD, Zhang X, Feng JT. Effect of allyl isothiocyanate on ultra-structure and the activities of four enzymes in adult *Sitophilus zeamais*. Pestic Biochem Physiol 2014; 109:12-17.
 44. Kimbaris AC, Kioulos E, Koliopoulos G, Polissiou MG, Michaelakis A. Coactivity of sulfide ingredients: a new perspective of the larvicidal activity of garlic essential oil against mosquitoes. Pest Manag Sci 2009; 65:249-254.