



E-ISSN: 2320-7078
P-ISSN: 2349-6800
JEZS 2017; 5(3): 1115-1120
© 2017 JEZS
Received: 06-03-2017
Accepted: 07-04-2017

Showket A Dar

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

GM Mir

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

GM Lone

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

Gh. I Hassan

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

Sajad H Parey

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

Correspondence**Showket A Dar**

Department of Entomology,
RTCPPPM-SKUAST-K,
Srinagar, Sheri-e-Kashmir
University of Agricultural
Science and Technology,
Shalimar, Jammu and Kashmir,
India

Potential habitat value for wild bees in Kashmir valley

Showket A Dar, GM Mir, GM Lone, Gh. I Hassan and Sajad H Parey

Abstract

The decrease in honeybee population in India has highlighted the importance of native bee conservation, and the need for research on the ecological requirements of native bees in various landscapes. In this study, we investigated the value of hedgerows as foraging habitat for native bees in mosaics of small scale agriculture and natural vegetation in the landscapes of Kashmir valley. In the spring of 2013-14, we surveyed bees and flowers in four habitats, i.e. hedgerows, agricultural fields, grasslands, and native woodland. We investigated that how hedgerows varied compared to other available habitats in bee abundance and species richness; the species composition in hedgerows compared to species composition in agricultural fields and woodland. We found that hedgerows were attractive foraging habitat for native bees, especially in early spring and summer. Cumulative species richness was highest in hedgerows, although cumulative species richness did not significantly differ among fields, hedgerows, and woodland. While bee faunas overlapped among habitats, and the bee assemblages in hedgerows were more similar to those in fields than to those woodlands. The flowering shrubs were important in attracting bees that were otherwise uncommon in the landscape, including some species that are potentially valuable pollinators of peach, plum and cherry.

Keywords: Hedgerows, habitat, bees, wild, ecology

1. Introduction

Recently UN science report warns of bees which are in the period of decline and there are going to be increasing consequences of their loss [19]. Habitat loss and fragmentation in agricultural landscapes lead to severe declines of abundance and richness of many insect species in the isolated semi-natural habitats [13]. The problem of habitat loss has been characterised as a “pollinator health crisis” where health refers to pollinator species diversity and abundance [7, 15]; and the pollination from these bees is most valuable ecological service provided by wildlife [8]. Agricultural landscapes are increasingly important settings for biological conservation, especially for the conservation of important pollinators such as native bees of Helictidae family [9, 10]. Bees (Hymenoptera: Apoidea) are the dominant providers of pollination services in both natural and agricultural ecosystems [2]. Field research with a range of crops suggests that maintaining the abundant and diverse native bee communities can provide insurance against the loss of pollination services in the face of reduced honeybee populations [26]. The potential of native bees to act as an alternative pollinators for crops is one reason to focus on their conservation near the farmland and other landscapes. Effective conservation of these native bee populations depends on understanding how the ecological requirements of wild bees can be met in agricultural landscapes [9]. Agricultural land can be insufficient to provide all of the resources necessary to sustain resident bee populations, e.g., pollen, nectar, floral oils, nest sites, nesting materials, and stable sites for over-wintering [11]. Intensively managed agricultural landscapes tend to lack species-rich floras and continuity of floral resources. The urban landscape offers a potential habitat or refuge to many different species [21]. Wild bees are amongst those seeking a safe haven in our back gardens, roadside verges, industrial sites and hedgerows. It was found that between 13-40% of wild bee species were found living in urban settings [3]. Barrier effects of hedgerows are also known to restrict movement of some bee species [14]. One means of preserving resources for wild bees in agricultural landscapes is maintaining patches of semi-natural vegetation within the agricultural matrix [24] for pollination as bees have complex habitat requirements. Not only do some bees have very specific demands for certain floral resources, they also require specific structures to build their nests. Hedgerows are important as for as construction of nesting structures are concerned. Pollen sources are often distributed over different habitats [26].

Therefore, most bee species rely on several partial habitats to cover all needs, and these habitats have to lie within the flight range of the bee species, which is a usually limited [6] and demand the hedgerows close to the agricultural areas. Here, we investigate the use of hedgerows by wild bee communities in mixed landscapes of native vegetation and small-scale agriculture in Kashmir. The study areas are two corridors in south-eastern Kashmir, noted for their high diversity of bees and flowering plants. Hedgerows are common features on the small farms of both study areas.

2. Materials and methods

The present investigations were carried out from 2013 to 2014 in different landscapes of Kashmir valley.

2.1 Study area

Kashmir division of State Jammu and Kashmir is located in

North-eastern of India. Geographically it is stretched between 32° 17' to 37° 60' N latitude and 73° 26' to 80° 30' E longitude. The mountain range in the Himalayas region varies in altitude with 5,550m on North-east dip down to about 2,770m on South. Generally, the Kashmir contains the upper stages of the forest vegetation including pinus, populus, willow, rubenia and some other social forestry trees and lower stages of agricultural and horticultural crops including apple, pear, peach, plum, apricot, almond and cherry. The research was conducted in Budgam, Pulwama and Srinagar situated at the heights of 5400, 1570 and 1645 meters from mean sea level (MSL). Higher altitudes of Kashmir (India) region are the centres of diversity, especially Budgam and Pulwama (Fig. 1), support diverse invertebrate faunas and species-rich floras. The two drainages have almost very similar plant communities. The dry uplands support diverse types of desert vegetation community.




Fig 1: Experimental locations were survey and samplings of wild bee species were conducted

This region of N-S receives 25-35% of its precipitation during the winter and early spring months, and 50-60% in the monsoon period. The agricultural landscapes of both the upper experimental plots of both Budgam and Pulwama districts and lower orchards of Srinagar are characterized by relatively small fields, less than 3 ha, and a mosaic pattern of habitat types. Fields are frequently separated by hedgerows, having abundance of native shrubs, herbs and mulberry plantations and other social forestry plants.

The matrix of field and hedgerow is interspersed with areas where various forest trees have been allowed to regenerate. In the areas of Kashmir Valley, agricultural land occurs in two distinct clusters. We surveyed hedgerows and other agricultural habitats on 10 locations, five along the south and five on the North (Fig: 1). The agricultural clusters of the study areas are largely made up of contiguous farms. In order to minimize any effect of farm-to-farm variation, we standardized farms for available habitat types and management practices -including crops grown and pesticide/herbicide use. Fields were planted with crops like maize, oats and brassica; however, large areas in all experimental sites are under paddy and considerable area is maintained as irrigated pastures of mixed grasses and forbs for cattle. On each farm, various types of social forest trees were used to graze cattle for at least part of the year. There was no aerial application of pesticide or herbicide on any farm; spot-application of herbicide was used to suppress woody seedlings in fields. In all experimental sites, the apple orchards were also surveyed for the bee species richness and

abundance.

We surveyed bees and flowers in four habitat types and established 15 transects in each category of landscape. Each transect consisted of a 2 m by 40 m rectangular plot, the maximum length of transects being constrained by the length of hedgerows on some farms. We placed transects randomly along hedgerows and placed a second transect in the approximate centre of the adjacent field, running parallel to the longest dimension of the field. We placed a third transect in a randomly chosen site in the nearest woodlot. The minimum distance between hedgerow and woodlot transects was 100 m. The minimum distance between hedgerow and field transects was somewhat smaller (75 m). We identified potential native woodland sites based on the presence of the dominant native perennial grass and/or the absence of economic shrubs. For practicality of sampling, once one transect was placed randomly within suitable woodland habitat, a second transect was placed 200 m away. This was the minimum distance among native woodland transects. Woodland transects were located between 400 and 600 m from the nearest agricultural field.

2.2 Bee sample collection

We used timed periods of observation and hand-netting to sample bee assemblages. Surveys were completed between 9:30 am and 4:00 pm, and were only conducted in clear weather. Afternoon surveys were also done but usually not feasible because of reliable afternoon cloud cover and rain in Kashmir during the monsoon season of 2013-14. Depending

on the survey year, two or three observers were used. Observers alternated the transects they surveyed, and the order in which transects at a given site were surveyed, in subsequent sampling periods. In each survey, the observer spent 30 min in the 20-35 m transect area, catching insect specimens and recording the flower species, if any, visited by each bee. Handling time was not included in the observation time; the clock was turned off when a bee was caught and while it was being processed. The 30 min total observation time was divided into five 6-min periods (one for each 7 m subplot) in order to spread observer attention across the transect area. Data from the four subplots were combined to make one sample per transect. We attempted to catch every bee detected, even if not caught photographs were taken while foraging. Voucher specimens were deposited in pollinator laboratory of entomology, SKUAST-K. The three sampling periods corresponded to the pre-monsoon, early monsoon, and late monsoon seasons. Flowering plant phonologies were surveyed from the lowest elevation sites in each round. The highest elevations were visited last within each round.

2.3 Data analysis

2.3.1 Bee species richness

In order to compare species richness among habitats with almost equal sampling intensity, we estimated total species richness for each habitat using ManiTab and O.P. Sheoran software, to determine the significance among the species pertaining to the respective specific orders. The diversity and abundance of the bees captured from each habitat and comparisons between particular pairs of habitats were made using the Student’s t-test. We compared mean species richness and abundance among habitats using one-way ANOVA. Separate ANOVA were performed for each sampling period, so that statistical comparisons of species richness and abundance were only made among observations made in the same sampling periods. We chose the Bray–Curtis index to quantify dissimilarity among bee assemblages.

3. Results

3.1 Total species richness

Altogether, we collected 531 individuals of 20 bee species in 9 genera and five families (Table 1). Sampling intensity varied among habitats, with a range of n = 65, 135 samples (one sample equals one survey of one transect during one sampling period in 1 year). Therefore, to compare species richness among habitats, we randomly conducted 65 surveys, and the analysis of the rough data collected yielded a total of 16 species in fields, 18 species in hedgerows, 20 species in fruit orchards near hedge rows, 16 species in native woodland, and 13 species in woodlots. Overall, 90% of the species recorded were common across different landscapes. In addition all of the species were recorded near by the stone fruit orchards. When cumulative species richness was plotted against number of surveys in each habitat, hedge rows near orchards had the highest estimated species richness for a given level of sampling effort. Estimated species richness in fields, hedgerows, and native wood land differ significantly. Estimated species richness in woodlots was significantly lower than in the other three habitats. Since, the flowering commences later at higher elevation, abundances of workers and males are also shifted later; therefore elevational comparisons play an important role in species richness [6]. While observed species richness was highest in hedge rows near orchards, further the jack-knife estimates of actual species richness were highest for hedgerows near orchards.

During the present investigations the *Lasioglossum* genus was the most abundant flower visitor, representing 45 to 48% of all individuals collected. Foragers of this species recorded at the study sites belong to wild colonies. The *Lasioglossum marginatum* was the most abundant species (Table.1) and foraging apparently takes place within a radius of 100 m from the nesting site. On stone fruit flowers in orchards of experimental locations, the per cent relative abundance of insect pollinators species were significantly highest of 5.0 ± 0.67 in 2013 ($P \leq 4.87$, t. stat. = 2.79, p.value < 0.01) and comparatively lowest 3.0 ± 0.35 pollinators/m²/10 min. in 2014 ($P \leq 9.75$, t. stat= 2.04, p.value < 0.01) with pooled relative abundance of 4.0 ± 0.51 pollinators/m²/10 min. (F. ratio 0.94; CV, 17.84; SE, 0.87; CD_(0.05) = 0.49; Pearson’s correlation = 0.90, T-test = 4.73, p.value < 0.001). The average relative abundance of Halictidae bees visiting the peach flowers was statistically significant and were in the order of abundance, *Lasioglossum margintaum*, *L. nursei*, *L. himalayans*, *L. regolatum*, *L. sublaterale*, *L. leucozoni* and *L. polyctor* with 4.0 ± 0.51 , 3.54 ± 0.34 , 3.27 ± 0.41 , 3.0 ± 0.21 , 1.5 ± 0.18 , 1.5 ± 0.31 and 1.89 ± 0.10 pollinators/m²/10 min, respectively. However, the relative abundance of *Halictus constructus* from genus *Halictus* of family Halictidae was comparatively minimum only of 2.28 ± 0.11 pollinators/m²/10 min. The species viz, *Andrena patella*, *A.flordula*, *A.cineraria* and *A. Bicolour* of andrenadae have relative abundance of 1.78 ± 0.14 , 1.56 ± 0.16 , 0.33 ± 0.09 and 0.89 ± 0.09 pollinators/m²/10 min, respectively (Table 1). In family Apidae the species *Xylocopa valga* and *X.violaceae* have the abundance of 1.17 ± 0.19 and 1.04 ± 0.11 pollinators/sq. meter/10 min, respectively. Among the family Megachilidae only two species *Megachile rotundata* and *Anthedium consolatum* were reported having the significant relative abundance of 0.95 ± 0.06 and 0.94 ± 0.39 pollinators/m²/10 min, respectively.

Table 1: Relative abundance of insect pollinators of stone fruit crops in Kashmir valley.

S. No	Pollinator Species	2013	2014	Pooled	SE
1	<i>Lasioglossum marginatum</i>	5.00	3.00	4.00	±0.51
2	<i>L. regolatum</i>	4.00	2.00	3.00	±0.34
3	<i>L. himalayense</i>	4.33	2.22	3.27	±0.41
4	<i>L. sublaterale</i>	1.67	1.33	1.50	±0.21
5	<i>L. leucozonium</i>	1.67	1.33	1.50	±0.18
6	<i>L. nursei</i>	4.78	2.33	3.54	±0.31
7	<i>L. polyctor</i>	2.00	1.78	1.89	±0.10
8	<i>Halictus constructus</i>	1.33	1.00	1.16	±0.11
9	<i>Sphecodes tantalus</i>	0.67	0.33	0.50	±0.09
10	<i>Andrena patella</i>	2.00	1.56	1.78	±0.14
11	<i>A. flordula</i>	2.34	0.78	1.56	±0.16
12	<i>A. cineraria</i>	0.44	0.22	0.33	±0.09
13	<i>A. bicolor</i>	1.35	0.44	0.89	±0.09
14	<i>A. barbilabris</i>	1.00	0.22	0.61	±0.19
15	<i>Amegilla cingulate</i>	0.45	0.33	0.39	±0.11
16	<i>Megachile rotundata</i>	1.11	0.78	0.95	±0.06
17	<i>Anthedium consolatum</i>	0.89	1.00	0.94	±0.39
18	<i>Xylocopa valga</i>	1.33	1.00	1.17	±0.19
19	<i>X. violacea</i>	1.07	1.00	1.04	±0.11
20	<i>Bombus sp.</i>	0.11	0.00	0.05	±0.01
	Total species =20	342	189	-	-
	N (Total samples collected=731), F. ratio 0.94; CV, 17.84; SE, 0.87; CD _(0.05) = 0.49; Pearson’s correlation = 0.90, T-test = 4.73, p. value < 0.001				

Table 2: One way ANOVA of species abundance across the two years of study.

F-Test Two-Sample for Variances		
	2013	2014
Mean	1.712632	1.034211
Variance	1.739343	0.494437
Observations	19	19
df	18	18
F.value	3.517826	-
P(F<=f) one-tail	0.005342	-
F Critical one-tail	2.217197	-

3.2 Species composition and relative abundance

There was broad overlap in species composition in the cumulative bee faunas of fields, hedgerows (orchards), woodlots, and native woodland. The majority of the hedgerow bee fauna was shared either with other on-farm habitats or with native woodland. Excluding singletons and using data only from 2013 to 2014 field seasons, 61% of species in hedgerows also occurred in fields, 48% also occurred in woodlots, and 71% were found in native woodland. Only 7.33% were found exclusively in hedgerows. Likewise, fields, woodlots, and woodland shared the majority of the species present there with other habitats- and had small proportions of 'unique' species. We also compared bee assemblages at the transect level- summing the abundances of bee species occurring on a given transect in the pre, early, and late monsoon over 2 years. Not surprisingly, the strongest difference was between fields and each of the hedgerows (orchards), woodlots and native woodland. Hedgerows were significantly more similar to woodlots and native woodland than to assemblages in agricultural fields. In general, there was high variability in bee assemblage composition among transects within habitats and wide overlap in composition among habitats. This variability was reflected in the low effect size scores for differences among habitats.

3.3 Spatial autocorrelation in bee species composition

From the dataset of cumulative bee assemblages, summing bee abundance data for each transect across all sampling periods in 2013 and 2014, to test for one way ANOVA in bee species composition (Table.2). Sample sizes in this dataset are lower than the total number of transects surveyed, as only those transects that had been in four of six sampling periods, and a minimum of three species in their cumulative bee assemblages, were included in the analysis. T-test showed a statistically significant (t -test < 0.05%), though relatively weak. This result was apparently primarily due to correlation among native woodland transects along the Pulwama, Budgam and Srinagar districts.

3.4 Abundance and species richness of bees by sampling period

Bee abundance was highly variable among transects within habitats, and patterns of abundance were variable across years. There were some consistent patterns, however. In each year, bee abundance in hedgerows peaked in the pre-monsoon, declined in the early monsoon, and then increased slightly in the late monsoon. During the pre-monsoon period, fields and hedgerows tended to have higher abundance than either native woodland or woodlots. Bee species richness was also variable within habitats, and there were few significant differences among habitats.

Pre-monsoon species richness was highest in hedgerows in 2013. This same trend was apparent, but not statistically

significant, in 2014. During the year 2014, the valley Kashmir was hit by floods and various foraging habitats were affected so bee assemblage and abundance were low. Fields tended to have the most species per transect during the early monsoon, a time when bee abundance and species richness was depressed in most habitats. Woodlots had the lowest mean species richness in most sampling periods. Pollen specialists foraged in each of the habitat types we surveyed, but habitats varied in specialist species richness. Cumulatively, hedgerows attracted a relatively large number of specialists – more than both fields and native woodland. With equivalent sampling effort, hedge rows had the highest number of specialist species as in order; hedgerows > fields > native woodlands > woodlots.

4. Discussion

Anthropogenic landscape elements, such as roadsides, hedgerows, edges, and power line clearings, can be managed to provide important habitats for wild bees. Hedge rows acted as net exporters of bees into adjacent fields [23] and also had enough bee populations. Within-farm habitat restoration such as hedgerow creation may be essential for enhancing the native pollinator abundance and diversity, and for pollination services to adjacent crops [17]. Semi-natural grasslands, such as calcareous grasslands, provide important habitats for bees [18], but are often lost due to changes in land use, particularly reduced livestock grazing [22]. However, other anthropogenic landscape elements, such as power line clearings [20], hedgerows [17], and orchard field edges [23], may also provide important habitats for bees in the agricultural landscape matrix. In the mixed farm and natural landscapes of our study areas along the three experimental sites, it is clear that hedgerows contribute to available foraging habitat for local native bee populations. Net surveys showed that diverse assemblages of bees are finding nectar and pollen resources in hedgerows; a total of 20 bee species from 9 genera and five families were observed visiting flowers in hedgerow transects. Because of the extreme temporal variability in native bee faunas, it is important to look at habitat use by season, not just in sum. In our study areas, hedgerows were arguably the best foraging habitat for bees. Hedgerows tended to attract more diverse bee assemblages than fields, woodlots or native woodland during the 2013-14. The flower resources of the dominant tree and native woodland-associated shrubs drove this pattern, attracting the majority of native bees foraging in hedgerows in pre-monsoon. The trend was apparent in both years of the study, but only statistically significant in 1 year; the non-significant difference in 2014 may have been due to underlying low variability in species richness among transects in both hedgerows and fields. Hedgerows bloom and so provide food for insects [16]. Research has shown that native wild plants, shrubs and trees within hedgerows provide important foraging resources for wild bees and managed honey bees [16, 17]. Both agricultural fields and hedgerows were better foraging habitat, in terms of bee abundance and species richness, than either woodlots or native woodland during the pre-monsoon. Fields attracted bees in higher or comparable numbers to hedgerows during the pre-monsoon in 2013 year, and in later sampling periods. These findings differ somewhat from the European literature, in that hedges and fields were equally attractive to wild bees for at least some seasons. Croxton [5] found that bumblebees in agricultural habitats find much higher numbers foraging in the herbaceous under-stories of hedgerows than in adjacent fields. While as, pasture and hay crop fields in our study sites, commonly supported perennial and annual flowering weeds in addition to

grasses and hay crops. Clearly, irrigation extends the period of available forage in fields beyond that of natural habitats, where available flower resources are tied to seasonal rains. This expanded blooming period could boost the total bee species richness observed in fields. In a study of bee communities in a landscape of forested heath, agriculture, and suburban development in New Jersey, Winfree [25] also found greater total species richness in agricultural habitats than in native forest. However, the pattern of bee species richness was found opposite to the one we observed. In most of the early and late flowering periods, the proportion of woodland transects with no flowers was relatively high.

Healthy hedgerows are home to a rich plant community, and provide crucial bee habitat [16]. For example, hedgerow shrubs such as cherries, plants of family rosacea, dogwoods, hawthorns and wild apple trees are a reliable and plentiful source of nectar and pollen in May and June, a time of year when many other plants have not yet flowered. For the most part, hedgerows appeared to offer additional resources for native bee species that were also using other agricultural and natural habitats in the landscape. Excluding singletons, hedgerows shared 87% of their bee species with at least one other habitat. The dispersed pattern of bee species distribution among hedgerows and other available habitats was evident in most sampling periods and in the cumulative bee faunas in each habitat. Winfree [26] found a similar pattern of wide overlap in the bee faunas of agriculture and native forest in their New Jersey study area. As in our data, a relatively small proportion of species occurring in agricultural or forest habitat was unique to one habitat type [26]. One explanation for the wide overlap in bee faunas among habitats in our study areas is the close proximity of multiple habitat types in the landscapes we studied. Depending on the bee species studied and methodology used, typical bee foraging distances are estimated at 150 m to more than 1.5 km. In our study areas, multiple agricultural and natural habitats are often available within a radius of 500 m to 1 km, well within the flight ranges of many native bee species. The generalized foraging behaviour of many bee species may also explain their wide distribution among available habitats on and off farms. Solitary bees commonly use multiple habitats to gather the resources they require, commuting between nest sites and foraging habitats, and tracking patchy, ephemeral floral resources [24, 2].

In intensive industrially farmed fields there was low field diversity (lack of hedgerows), and bee abundance and diversity were lowest [1]. The food resources in hedgerows attract some native bee species that were otherwise uncommon in the landscape, and even some species are unique in their preference to floral resources in hedgerows. The most abundant examples were *L. marginatum*, *L. nursi* and *L. himalayans*. Hedgerows attracted these native bees in significantly greater numbers and with significantly greater frequency than either agricultural fields or woodland. Andrenidae species were only rarely found in habitat other than hedgerows, and *A. cinera* was only caught in hedgerows. Helictidae species preferentially visited the flowers of a native shrub, fruit plants and agricultural crops; however present survey was done in landscapes near fruit orchards. Hedgerows may indirectly contribute to local bee diversity by providing forage to an assemblage of native bee species that vary widely in foraging ecology and seasonal activity period. Further the bee fauna in hedgerows included some species that are generalists in both habitat use and pollen collection (e.g., *Lasioglossum*, *Bombus*, and *Xylocopa* spp.).

5. Summary and conclusion

In intensive farmed fields there was low field diversity (lack of hedgerows) and bee abundance and the bee diversity were lowest. The most abundant examples species were *L. marginatum*, *L. nursi* and *L. himalayans* found near to the hedgerows. Hedgerows attracted these native bees in significantly greater numbers and with significantly greater frequency than either agricultural fields or woodland. Andrenidae species was only rarely found in habitat other than hedgerows, and *A. cinera* was only caught in hedgerows. In this study, we investigated the value of hedgerows as foraging habitat for native bees in mosaics of small scale agriculture and natural vegetation in the landscapes of Kashmir valley. Hedgerows varied significantly in bee abundance and species richness. We found that hedgerows were attractive foraging habitat for native bees, especially in early spring and summer. Cumulative species richness was highest in hedgerows, although cumulative species richness did not significantly differ among fields, hedgerows, and woodland. So in future we focus on the habitat conservation so as to meet the pollination needs of economic crops in the Kashmir valley.

6. Conflict of interest

There is no conflict of interest among the authors.

7. Acknowledgement

Authors are highly thankful to DST, New Delhi for financial assistance to conduct this research. Further, authors are also thankful to SKUAST-K, Srinagar for the laboratory and internet facilities provided during the studies

8. Reference

- Allsopp M, Tirado R, Johnston P, Santillo D, Lemmens P. Plan bee-living without pesticides moving towards ecological farming. Greenpeace International, Amsterdam, 2014, 21-39.
- Cane JH. Habitat fragmentation and native bees: a premature verdict? Conservation Ecology. 2001; 5(1):3.
- Colla SR, Ratti CM. Evidence for the decline of the western bumble bee (*Bombus occidentalis* Greene) in British Columbia. Pan-Pacific Entomologist. 2010; 86(2):32-3.
- Cornelissen ACM. 2012. Bijen in en rond de stad; een literatuurstudie. Plant Research International, Wageningen UR Postbus 69, 6700 AB, Wageningen. Entomologische Berichten, 2012; 72(1-2):120-124.
- Croxtton PJ, Carvell C, Mountford JO, Sparks TH. A comparison of green lanes and field margins as bumblebee habitat in an arable landscape. Biological Conservation. 2002; 107:365-374.
- Gathmann A, Tschardt T. Foraging ranges of solitary bees. Journal of Animal Ecology. 2002; 71:757-764.
- Goulson D, Nicholls E, Botias C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science 2015; 347(6):22-29).
- Hall A, Andrea B, Nicole S. Sustainability science for urban pollinator research and conservation. Ecology and Jesuit in communication. 2016, 1-6.
- Jauker F, Bondarenko B, Becker HC, Steffan-Dewenter I. Pollination efficiency of wild bees and hoverflies provided to oilseed rape. Agric. For. Entomol. 2012; 14:81-87.
- Klein AM, Vaissiere BE, Cane JH, Kremen C,

- Tscharntke T. Importance of pollinators in changing landscapes for world crops. *Proc R Soc Lond B*. 2007; 274:303-313.
11. Kremen C, Williams NM, Aizen MA, Gemmill-Herren B, LeBuhn G, Minckley RL *et al*. Pollination and other ecosystem services provided by mobile organisms: a conceptual framework for the effects of land- use change. *Ecol. Lett*. 2007; 10:299-314
 12. Krewenka K, Holzschuh A, Tscharntke T, Dormann CF. Landscape elements as potential barriers and corridors for bees, wasps and parasitoids. *Biological Conservation*. 2011; 144:1816-1825.
 13. Kuefler D, Hudgens B, Haddad NM, Morris WF, Thurgate N. The conflicting role of matrix habitats as conduits and barriers for dispersal. *Ecology*. 2010; 91:944-950.
 14. Maddox D. Can cities save bees? How can urban habitats be made to serve. The nature of cities. *Pol. Conservation? How can that story be better told?* 2016, 1-7
 15. Monkman D. Hedgerows are nature's ally in farm fields, 2013, 1-4
 16. Morandin LA, Kremen C. Hedgerow restoration promotes pollinator populations and exports native bees to adjacent fields. *Ecol Appl*. 2013; 23(4):829-39.
 17. Murray TE, Fitzpatrick U, Byrne A, Fealy R, Brown MJF, Paxton RJ. Local-scale factors structure wild bee communities in protected areas. *J Appl. Ecol*. 2012; 49:998-1008.
 18. Potts S. UN-sponsored scientific report warns too many species of pollinators are nearing extinction. *U.S. News and World Report*, 2016, 1-3.
 19. Russell K, Ikerd H, Droege S. The potential conservation value of un-mowed powerline strips for native bees. *Biol. Conserv*. 2005; 124:133-148.
 20. Samnegard U. The impact of forest on pest damage, pollinators, and pollination services in Ethiopian agricultural landscape. Thesis submitted to department of ecology, environment and plant science, Stockholm University, Ethiopia. 2016; 7(6):49-354.
 21. Stoate C, Báldi A, Beja P, Boatman N, Herzon I, Van Doorn A *et al*. Ecological impacts of early 21st century agricultural change in Europe—A review. *J Environ. Manage*. 2009; 91:22-46.
 22. Sydenham MAK, Eldegard K, Totland Q. Spatio-temporal variation in species assemblages in field edges: seasonally distinct responses of solitary bees to local habitat characteristics and landscape conditions. *Biodivers. Conserv*, 2014; 23:2393-2414.
 23. Westrich P. Habitat requirements of Central European bees and the problems of partial habitats. *The Conservation of Bees* (eds A. Matheson, S.L. Buchmann, C. O'Toole, P. Westrich & I. H. Williams), Academic Press, London, 1996, 1-16.
 24. Winfree R, Bartomeus I, Cariveau DP. Native pollinators in anthropogenic habitats. *Annu. Rev. Ecol. Evol. Syst*. 2011; 42:1-22
 25. Winfree R, Williams NM, Dushoff J, Kremen C. Native bees provide insurance against ongoing honey bee losses. *Ecology Letters*, 2007; 10:1105-1113.