



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2018; 6(6): 1354-1357

© 2018 JEZS

Received: 22-09-2018

Accepted: 26-10-2018

**SS Lonkar**Om Satya Sai College,  
Jawaharnagar, Bhandara,  
Maharashtra, India**RV Tijare**Department of Zoology,  
Institute of Science, Nagpur,  
Maharashtra, India**GT Kedar**Department of Zoology,  
Government of Maharashtra's  
Ismail Yusuf College, Jogeshwari  
(E), Mumbai, Maharashtra,  
India

## Studies on urban lakes of Nagpur, Maharashtra (I.) with special reference to Ichthyofaunal diversity

SS Lonkar, RV Tijare and GT Kedar

### Abstract

The present study carried at three lakes of Nagpur City in Central India from July 2012 to Jun 2014 confirmed Ichthyofaunal diversity of 21 species belonging to 9 different orders, *viz* Cypriniformes, Anguilliformes, Beloniformes, Paraformes, Singuilliformes, Clupeiformes, Mastacembaliformes, synganthiformes and Ophiocephalliformes. Maximum fish diversity was recorded in Telangkhedi lake (6.66) followed by Ambazari lake (5.33) and Gandhisagar lake (4.00). Presence of carps like Catla-catla, Labeo rohita, Cirrihina mrigala and Silver carp showed good productive grounds for fish culture practice in three lakes.

**Keywords:** Nagpur, lakes, fish fauna, fish culture

### Introduction

The Indian Subcontinent is endowed with a vast expanse of open inland fresh water bodies which includes a large number of rivers, ponds, dams, impoundments and lakes. These fresh water bodies provide source of bio-diversity and ecological assessment. It also provides a source of socio-economic development through various activities. Fishing is one of the ancient practices carried out since early period. Fish is a valuable source of protein and occupies a significant position in the socio-economical fabric of South Asian countries. For effective exploitation of any aquatic ecosystem basic information on its biodiversity is a must, thus there is a need to survey fish fauna associated with different freshwater habitats which will help in planning methods for their effective exploitation for fish production.

Many workers have studied taxonomy, bio-diversity and distribution of fishes found in various parts of Indian subcontinent provided that there is a need for the survey of bio diversity of fishes in different types of habitats all over the country. Jayaram (1981) [6] studied fish diversity of Indian subcontinent. Yazdani (1994) [18] reported Ichthyofauna from Krishna, Cauveri and Ganga River. In State of Maharashtra, Ichthyofaunal diversity was studied by Ahirrao and Mane (2000) [1], Sakhare & Joshi (2003) [10], Yadaw (2003) [16], Yadaw (2006) [17], Rathod *et al.* (2008) [9], Tijare and Thosar (2008) [14] and Harney *et al.* (2009) [5]. However, very less information is available about ichthyofauna present in lentic and lotic habitat of Nagpur district. Present study aims to document the fish fauna of three lakes of Nagpur city in Central India.

### Study Area

Nagpur city [21°07'N & 79°07'E] the second capital of Maharashtra state lies in centre of India. The city has dry subtropical monsoon climatic condition with temperature range 06 °C - 45 °C (Geo. Inf. Nat. Inf. Centre, 2006) (Figure 1). Based on old records, city was boasted of about twenty two water bodies, but rapid urbanization, load of population, widespread encroachment and continuous logging resulted into the existence of few water bodies in and around city. Primary aim of study is to investigate the ichthyofauna of three lakes namely Futala Lake, Gandhi Sagar lake and Ambazari Lake (Figure 2). As these lakes were used in many activities since time period, fishing is one of old practice. Fishing is carried out throughout the year and fish culture is done subsequently in respective three lakes.

### Futala Lake

[21°09'N. & 79.09'E] also known as Telangkhedi lake is an ancient and historical lake exists for 200 years and situated beside the highways on western area of Nagpur city.

### Correspondence

**SS Lonkar**Om Satya Sai College,  
Jawaharnagar, Bhandara,  
Maharashtra, India

The Futala Lake furnished with immense finery and magnificence is one of the must visit tourist attractions of the Nagpur city. Spread over an extensive are of 60 acres, the lake is enclosed by the guard walls on all the sides and encircled by a granite stone paving bequeathed with a fascinating garden.

#### Ambazari Lake

[21°10'N & 79°05'E] is almost a natural reservoir formed in the basin of the Nag river situated on the western outskirts of Nagpur. The lake covers an area of 15.4 km<sup>2</sup> having storage capacity of 1837 lakh gallons of water. It is the largest lake in the city.

#### Gandhisagar Lake

[21°08' N. & 79°08' E] is located near Raman Science Centre about one km. east of Nagpur. The lake which is said to be exists for more than 275 years established as a source of water supply by the then rulers. The picturesque rectangular shaped Gandhi Sagar reservoir is now enclosed with stonewalls and iron railings. One can also found a small island in the middle of the lake with an attractive Shiva temple and a garden.

#### Material and Methods

The present study was carried out in two years from June 2012 to July 2014. After netting the fishes, photographs were taken and the specimen were preserved in 10% formalin after giving abdominal cut and brought to laboratory for identification. For identification of fishes standard keys of Days (1878) [3], Jayaram (1981) [6] and Talwar and Jhigran (1991) were followed. Ichthyofaunal diversity of three lakes is reported in Table 1. Ichthyofaunal frequency, abundance and diversity was calculated by following statistical analysis methods of Zar (2005) [19].

Frequency (%) = T1/T2

T<sub>1</sub> = Total No. of pond in which species occur

T<sub>2</sub> = Total No. of ponds studied. Abundance = Frequency/100

Diversity of lake = Ts/T<sub>2</sub>

T<sub>s</sub> = Total No. of species found in lake

T<sub>2</sub> = Total No. of lakes studied.

#### Observation and Discussion

During the present investigation a total of 21 fish species belonging to 9 different orders were recorded from all the three lakes. The Futala Lake showed high species diversity with 20 species while Ambazari Lake showed 16 species, however 12 species were recorded from Gandhisagar Lake. Sharma *et al* (2011) [11] observed 15 fish species in Pinhole

Lake of Rajasthan.

Ten species of order Cypriniformes *viz.* Ctenopharyngodon idella (Grass carp), Hypophthalmichthys molitrix (Silver carp), Catla catla (Catla), Labeo rohita (Rohu), Cyprinids' carpio (Common carp), Cirrhina mrigala (Mrigal carp), Clarias batracus (Walking cat fish), Heteropneustus fossilis (Stinging cat fish), *Barbus* sp. (Barbel )and Wallago attu (Mully cat fish) dominated the lakes. Similar findings were also observed by Shinde *et al.* (2009) [12] in Harsool-Savangi dam. The Cypriniformes are an order of ray-finned fish including the carps and minnows and are most diverse in southeastern Asia (Nelson 2006) [8]. Cypriniform species are extremely variable morphologically and ecologically. The latter is evident from their wide distribution that includes virtually every type of freshwater habitat and an amazing diversity of reproductive and life-history strategies (Winfield & Nelson, 1991 and MacDonald 2008) [15, 7], Order Cypriniformes was followed by Ophiocephaliformes, Paraformes and Singuiliformes with 2 species while Anguiliformes, Beloniformes, Clupeiformes, Mastacembaliforms, syngnathiformes represented only 1 species respectively. Fish diversity was observed as maximum in Telangkhadi Lake (6.66) followed by Ambazari (5.33) and Gandhisagar Lake (4).

Presence of carps like *Catla catla*, *Labeo rohita*, *Cirrhina mrigala* and *Hypophthalmichthys molitrix* shows good productive grounds for fish culture practise in three lakes. Presence of *wallago Attu*, *Mystus seenghala* and *Clarius garripinnus* as cat fishes also provides embossing of healthy ecosystem in these lakes. The Ichthyofaunal diversity of Futala lake was dominant (6.66) followed by Ambazari lake (5.33) and Gandhisagar lake (4.00). Abundance is due to the easy availability of protein rich invertebrates and other food such as macrophytes, macrobenthic organisms and planktons. Every organism maintains specific relation with the environment in which it lives. These relations entail different environmental parameters eg. Temperature, humidity, diet requirements etc. (Blair RB 2001) [2]. The result of our survey highlight the fact that ichthyofauna in these lake is abundant which indicate the favorable condition for their survival.

However, Now-a-days these lakes are getting deteriorate by the activities of urban development, idol immersions, resultant stress and encroachments ultimately causing threat to fish fauna. Therefore, the conservation of these fragile ecosystems rich with diverse fish fauna is an essential and urgent task which can be achieved by reducing anthropogenic activities and introducing safe, environment friendly fish culture practice.




Fig 1: Map showing location of Nagpur city

Fig 2: Google map location of three lakes in Nagpur city

**Table 1:** Ichthyofaunal diversity of three lakes in Nagpur, (+ Present: - Absent)

Sr. No	Common Names	Scientific name	Order	Futala Lake	Ambazar i Lake	Gandhi-Sagar Lake
1	Grass carp	<i>Ctenopharyngdon idella</i>	Cypriniformes	+	+	+
2	Silver carp	<i>Hypophthalmichthys molitrix</i>	Cypriniformes	+	+	+
3	Catla	<i>Catla catla</i>	Cypriniformes	+	+	+
4	Rohu	<i>Labeo rohita</i>	Cypriniformes	+	+	+
5	Common carp	<i>Cyprinus carpio</i>	Cypriniformes	+	+	+
6	Mrigal carp	<i>Cirrhina mrigala</i>	Cypriniformes	+	—	+
7	Walking cat fish	<i>Clarias batracus</i>	Cypriniformes	+	+	—
8	Stinging cat fish	<i>Heteropneustus fossilis</i>	Cypriniformes	+	—	+
9	Barbel	<i>Barbus sp.</i>	Cypriniformes	+	+	—
10	Mully cat fish	<i>Wallago attu</i>	Cypriniformes	+	+	—
11	Indian mottled eel	<i>Anguilla bengalensis</i>	Angulliformes	+	—	—
12	Needle fish	<i>Belone cancila</i>	Beloniformes	+	—	—
13	Common snake headed	<i>Channa striatus</i>	Ophiocephalliformes	+	+	+
14	Spotted headed snake	<i>Channa punctatus</i>	Ophiocephalliformes	+	+	—
15	Bulls eye headed snake	<i>Channa nama</i>	Paraformes	+	+	—
16	Bulls eye gobby	<i>Glossogobius giuris</i>	Paraformes	+	+	+
17	Giant river catfish	<i>Mystus seenghala</i>	Singuliformes	+	+	+
18	Shrptooth catfish	<i>Clarius garipinnus</i>	Singuliformes	+	+	—
19	Clown knife fish	<i>Notopterus chitala</i>	Clupeiformes	—	—	+
20	Tire track eel	<i>Mastacembalus aramatus</i>	Mastacembaliformes	+	+	—
21	Mosabbique Tilapia	<i>Tilapia sp.</i>	Syngnathiformis	+	+	+

**Table 2:** Frequency and abundance of fish species

Sr. No	Common Names	Scientific name	Frequency	Abundance
1	Grass carp	<i>Ctenopharyngdon idella</i>	100	1
2	Silver carp	<i>Hypophthalmichthys molitrix</i>	100	1
3	Catla	<i>Catla catla</i>	100	1
4	Rohu	<i>Labeo rohita</i>	100	1
5	Common carp	<i>Cyprinus carpio</i>	100	1
6	Mrigal carp	<i>Cirrhina mrigala</i>	66	0.66
7	Walking cat fish	<i>Clarias batracus</i>	66	0.66
8	Stinging cat fish	<i>Heteropneustus fossilis</i>	66	0.66
9	Barbel	<i>Barbus sp.</i>	66	0.66
10	Mully cat fish	<i>Wallago attu</i>	66	0.66
11	Indian mottled eel	<i>Anguilla bengalensis</i>	33	0.33
12	Needle fish	<i>Belone cancila</i>	33	0.33
13	Common snake headed	<i>Channa striatus</i>	100	1
14	Spotted snake headed	<i>Channa punctatus</i>	66	0.66
15	Bulls eye snake headed	<i>Channa nama</i>	66	0.66
16	Bulls eye gobby	<i>Glossogobius giuris</i>	100	1
17	Giant river catfish	<i>Mystus seenghala</i>	100	1
18	Shrptooth catfish	<i>Clarius garipinnus</i>	66	0.66
19	Clown knife fish	<i>Notopterus chitala</i>	33	0.33
20	Tire track eel	<i>Mastacembalus aramatus</i>	66	0.66
21	Mosabbique Tilapia	<i>Tilapia sp.</i>	100	1

**Table 3:** Ichthyofaunal diversity of three lakes

Lakes	Diversity
Futala	6.66
Ambazari	5.33
Gandhisagar	4.00

## Conclusions

The study of ichthyofauna in the lakes of Nagpur reveals significant biodiversity, with a total of 21 fish species recorded across three lakes. Futala Lake exhibited the highest species diversity, followed by Ambazari and Gandhisagar lakes. The presence of key species such as carps and catfish highlights the lakes' potential for sustainable fish culture. However, urbanization, pollution, and encroachments are threatening these ecosystems. Conservation efforts focusing on reducing anthropogenic impact and promoting environmentally friendly fish culture practices are crucial to preserving the fish biodiversity and maintaining the

ecological balance of these water bodies.

## References

- Ahirrao SD, Mane AS. The diversity of Ichthyofauna taxonomy and fishes from freshwater bodies of Parbhani district, Maharashtra state (I). J Aqua Biol. 2000;15(1 & 2):40-43.
- Blair RB. Birds and butterflies along urban gradients in two eco-regions of the United States: Is urbanization a homogeneous fauna? In: Lockwood JL, McKinney ML, editors. Biotic homogenization: The loss of diversity through invasion and extinction. New York, Kluwer

- Academic Publishers; c2001, p. 33-56.
3. Days FS. The fishes of India. London: William Dawson and Sons Ltd.; c1878.
  4. National Informatics Centre. Geographical information on Nagpur city. Nagpur: National Informatics Centre; c2006. [Cited 2006 Jun 30]. Available from: [URL]
  5. Harney NV, Dhamani AA, Andrew RJ. Studies on Ichthyofaunal Diversity of three water bodies near Bhadrawati, Dist. Chandrapur MS (I). *Hislopia J.* 2009;2(2):151-157.
  6. Jayaram KC. The freshwater fishes of India, Pakistan, Burma and Sri Lanka. Handbook of Zoological Survey of India, No. 2. Calcutta: Zoological Survey of India; c1981, XII + 475 p.
  7. MacDonald M. Loaches: Natural History and Aquarium Care. [Hardcover]. Neptune City, NJ: TFH Publications, Ltd.; c2008.
  8. Nelson JS. Fishes of the World. 4th ed. Hoboken, NJ: John Wiley & Sons, Inc.; c2006. ISBN 0-471-25031-7.
  9. Rathod SD, Malu RA, Dhabade DS, Patil PS, Charjan AP, Wanjari HV. Diversity of fish fauna of Umra reservoir, Washim district, Maharashtra. *J Aqua Biol.* 2008;23(2):26-28.
  10. Sakhare VB, Joshi PK. Reservoir fishery potential of Parbhani district of Maharashtra. *Fishing Climes.* 2003;23(5):13-16.
  11. Sharma R, Sharma V, Sharma MS, Verma BK, Modi R, Gaur KS. Studies on limnological characteristics, planktonic diversity and fishes (species) in Lake Pichhola, Udaipur, Rajasthan (India). *Universal J Environ Res Technol.* 2011;1(3):274-285.
  12. Shinde SE, Pathan TS, Bhandare RY, Sonawane DL. Ichthyofaunal diversity of Harsool-Savangi dam, Dist. Aurangabad (M.S.) India. *World J Fish Mar Sci.* 2009;1(3):141-143.
  13. Talwar PK, Jhingran AG. Inland fishes of India and adjacent countries. Vol. 1 and 2. New Delhi: Oxford and IBH Pub Co. Ltd.; c1991.
  14. Tijare RV, Thosar MR. Ichthyofaunal study from the lakes of Gadchiroli District, Maharashtra (I). *J Aqua Biol.* 2008;23(2):29-31.
  15. Winfield IJ, Nelson JS. Cyprinid fishes: systematics, biology and exploitation. London: Chapman and Hall; c1991.
  16. Yadaw BE. Ichthyofauna of the northern part of Western Ghats. *Rec Zool Surv India.* 2003;Occ Paper 215:1-40.
  17. Yadaw BE. Pisces fauna of Tadoba Andhari Tiger Project conservation area. *Sec.* 2006;25:137-160.
  18. Yazdani GM. Biodiversity of fishes of river Ganga. *Report Zool Surv India.* 1994;72.
  19. Zar JH. Biostatistical Analysis. 4<sup>th</sup> Ed. Upper Saddle River, NJ: Pearson Education Inc.; c2005.