



E-ISSN: 2320-7078

P-ISSN: 2349-6800

www.entomoljournal.com

JEZS 2021; 9(6): 14-19

© 2021 JEZS

Received: 18-08-2021

Accepted: 26-09-2021

Ghazwan T KhudairMosul Directorate of Education
Affairs, Ministry of Education,
Mosul, Iraq**Atallah F Mekhlif**Department of Biology, College
of Education for Pure Science,
Mosul University, Mosul, Iraq

Antibacterial activity of secretion/excretion blow fly, *Callifora vomitoria* (Diptera: Calliphoridae) third instar larvae *in vitro*

Ghazwan T Khudair and Atallah F Mekhlif

Abstract

The antibacterial activity of excretion/secretion (ES) from flesh feeding blow fly, *Calliphora vomitoria* L. were relatively compared with that of Ceftriaxone (CRO) as standard drug. The larval ES was obtained from both normal fed larvae and others cannibalized infected larvae with the four studied pathogenic bacteria. Diffusion disc test was used to assess ES viability against the sensitive and piperacillin resistant bacterial species; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. The antibacterial potency of the two fractionation phases of ES in DMSO were varied among that four species. This potency depends upon the fractionated ES phase. Generally, dense (upper) and clear (lower) phases of normal ES more effective against Gram-positive than Gram-negative species, while not that for phases of ES collected from cannibalized infected larvae. It was found the growth inhibition zones significantly very varied between antibacterial activity of ES and CRO. Overall, the present result gives opportunity promise for qualitative evolution new resources against antimicrobe from larvae products.

Keywords: *Calliphora vomitoria*, excretion, secretion, antibacterial, resistant bacteria

Introduction

Carriens and corpus are suitable sources for nutrition and oviposition sites to some decomposing insect species, In addition innate immunity system, which represented by humoral and cellular immunity and that including melanization, phagocytosis, nodulation and encapsulation^[14] and^[24], some insects were supplied with external defense equipment characterized by antimicrobial activity, for instances oral and anal secretions of carrion beetles inhibited growth Gram-positive and Gram-negative bacteria^[11] and^[9]. Many dipteran larvae are exposed to stress of pathogenic microorganisms, so, larval excretions/secretions giving optimal activity against wide-spectrum of bacteria^[12] and^[7]. *Lucilia* and *Calliphora* are important genera in calliphorid family, and they depending in forensic entomology^[4] and^[5], these larvae have selective adaptation in contaminated environment with microbes, so, protected themselves by antimicrobial peptides (AMPs) which done separately and in synergistic each other^[21] and^[6], or potentiation by enhancing antimicrobial activity of others^[28]. Now these AMPs are be considered strong natural antibacterial inhibitors^[29]. Also, the extracted lipid compounds from *Calliphora vicina* and *Calliphora vomitoria* had antimicrobial activity^[8]. Excretion/secretion (SE) of *Lucilia sericata* larvae are use in healing necrotic wounds. ES were prevented biofilm formation and degraded biofilms initiation by the bacteria *Staphylococcus aureus* and *Pseudomonas aeruginosa*^[27].

First attention was paid to the beneficial effect of larval ES early in 1930^[23] was found that after 5 to 10 min – exposure, significant antimicrobial activity of non - sterile larvae of *L. sericata* against some pyogenic bacteria. The insects have ability to resist microbial infection^[15]. Injured *L. sericata* larvae were stimulated to produce lucifencin AMP with antibacterial activity^[26]. AMP are mainly produced by fat bodies and hemocytes and released to hemolymph^[13] and^[30]. AMPs are antibacterial molecules and important ingredients in the ES of the larval stage, besides to digestive enzymes which used in external digestion^[19]. The antibacterial molecules have been identified whether their sources SE or body extract^[5].

Since the scientific research signals application the hypothesis, carrion insects are protect themselves against microbial infection by antimicrobial molecules. In this scope, the present study aims to test the antibacterial activity SE of late instar larvae *Calliphore vomitoria* on

Corresponding Author:**Atallah F Mekhlif**Department of Biology, College
of Education for Pure Science,
Mosul University, Mosul, Iraq

growth inhibition the pathogenic bacteria *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumonia* *in vitro*.

Materials and Methods

Materials

Insect colony

Out door and nearby Entomology laboratory, the colony of the blue bottle fly, *Calliphora vomitoria* was established in semi-field condition. The cage dimensions were 2×1×1 m with wooden floor and sheltered for the colony maintenance, wild adult flies by net were collected from city slaughter in November 2020 and fed on 15% honey bee solution, fish remaining filth and ovaries were presented from local fishing market and used for larval feeding and ovipositing sites. Wet sandy soil was placed in cage beside feeding site for pupation the wandering full grown larvae.

Bacteria

The bacteria were used as marker to evaluate the antimicrobial activity of larval excretion/secretion. The pathogenic bacteria; *Staphylococcus aureus* (Gram-positive), *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumonia* (Gram-negatives) were obtained as a loan from

Dr. Alaa Taha Younis, Microbiology lab / Research unit / Biology Department / Collage of Education for Pure Science / Mosul University /Iraq.

Culture media

From NEOGEN Culture Media, Muller-Hinton media were purchased and culture prepared for bacteria inoculation

Methods

Preparation of excretion / secretion

The excretion/secretion (ES) were collected according to [13] and [18] with modifications. Approximately 1000 larvae (700 gm) in the late third stage were picked out from caged colony (plate 1). The larvae were preserved in enamel dish (25 × 25×20 cm), covered and tied with mesh clothe and left for overnight under the laboratory conditions. The larvae were rinsed by 100 ml of the negative solvent DMSO with 7: 1 (w/v) ratio. After removing the larvae, the obtaining crude were filtrated with wattmann filter paper No. 1 under low pressure. The filtrate was centrifuged at 3000 rpm for 15 min., and fractionated into two phases, they can be physically distinguished as upper deep and lower pale pink color (plate 1E). With about ration 20: 80 between the two rested phases.



Plate 1: Experimental materials of *Calliphora vomitoria* for preparing the excretion and secretion (SE); A – Ova, B – feeding larvae, C – collected larvae, D – larval cannibalism on infected larvae and E – ES filtrated phases.

Excretion and secretion of infected larvae

Infected *C. vomitoria* larvae were obtained by feeding them on another dead larvae as fellow; 150 larvae at second instar were took from the colony and fed on broth agar media contaminated with mixture of the four studied marker bacteria; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. Later, the grown third instar larvae were killed by cooling, and introduced as food to the developing experimental 150 second instar larvae plate 1C). The feeding larvae on dead others were grew and became in third instar. Then, same protocol to prepare excretion / secretion of uninfected larvae has been followed.

Antibacterial test

Bacteria growth inhibition by third larvae ES were *in vitro* tested on the marker bacteria, by agar diffusion discs, the test was modified after Kirby-Bauer [2], to estimate bacteria susceptibility. From each marker bacteria plate, samples were swapped on the petri dish plates, on each plate positive and negative controls and treated discs were fixed with three replicates, the discs were previously treated with three concentrations of ES were 0.05, 0.1 and mg/ml of each the two phases of the ES stock solution, and dissolved in 1.0 ml of the inert DMSO solvent. The plates were incubated for 24 hrs. at 37°C, the diameter of growth inhibition zones were measured by caliper in mm, and exclusive 5 mm (diameter of

the disc) of the readings. For evaluating the antibacterial activity of the ES, the growth inhibition zones were ranked after [26]. into the following grades: ≥ 8 mm good, 6-8 mm moderate, 4-5 mm and 2-3 mm very weak.

Results

Antibacterial activity of excretion / secretion (ES) of blue-bottle *Calliphora vomitoria* were studied against sensitive and resistant Gram-Positive *Staphylococcus aureus* and Gram – negatives; *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. The two main sources of the investigated ES were obtained from uninfected larvae and others infected with same experimental bacteria. Table 1 illustrates that the inhibition zone for sensitive bacteria resulting by the two supernatant phases of ES normal fed larvae in comparison with CRO antibiotic as positive control. The clear (lower) phase have significant growth inhibition

activity among the treated bacteria, the diameters of the inhibited zones were 21.5, 5.5 17.2 and 13.3 mm for the Gram - positive bacteria *S. aureus*, and Gram – negatives *E. coli*, *P. aeruginosa* and *K. Pneumoniae* respectively. Antibacterial activity of the dense (upper) phase less than that of the clear phase, and these inhibition zones with the same bacterial sequence were: 17.2, 12.8, 11.8 and 9.9 mm. On the other hand, ES of the cannibalized infected larvae were showed in Table 2, their inhibition zones 28.3, 21.8, 336.2 and 17.3 mm at lower SE phase, while inhibitions zones of the upper phase were 25.0, 17.2, 23.2 and 12.8 mm for the bacteria *S. aureus*, *E. coli*, *P. aeruginosa* and *K. Pneumoniae* respectively. Inhibition growth zones by CRO antibiotic were less than zone diameter of the two phase of infected or normal fed larvae (9.2, 3.3, 6.8 and 3.5 mm) for *S. aureus*, *E. coli*, *P. aeruginosa* and *K. Pneumoniae* bacteria.



Plate 2: Growth inhibition zones of *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella pneumoniae* and *Escherichia coli* were treated by clear (C) and dense (D) SE fractions of *Calliphora vomitoria* larvae. CRO – Ceftriaxone.

Antibacterial activity against resistant bacteria

Inhibition growth zones of the resistant bacteria *S. aureus*, *E. coli*, *P. aeruginosa* and *K. Pneumoniae* to piperacillin antibiotic were measured after treatment with two concentrations of each dense and clear phases of *C. vomitoria* ES. Table 3 shows the bacterial inhibition at the concentrations 0.1 and 0.05 mg/ml at clear (lower) phase for the resistant bacteria *S. aureus*, *E. coli*, *P. aeruginosa* and *K. Pneumoniae* were (13.8, 10.0 mm), (9.4, 5.5 mm), (16.8, 10.7

mm) and (14.0, 3.5mm) respectively at the concentrations. Besides, the clear (lower) phase of the ES were continued in antibacterial action but not proportionally with Concentration multiplication, as in dense phase application, so the inhibition growth zones; (15.2, 9.2 mm), (11.4, 9.9 mm), (12.8, 9.8mm) and (9.5, 4.2 mm) of the resistant bacteria (table 2). The +ve control (CRO antibiotic) had negligible effect (plate 3) on the treated resistant bacteria *S. aureus*, *E. coli*, *P. aeruginosa* and *K. Pneumoniae*.




Plate 3: Antibacterial activity of two SE fractionation (clear and dense) phases of *Calliphora vomitoria* on Piperacillin resistant bacteria *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella pneumoniae* and *Escherichia coli*. C – clear, D – dense phases. CRO – Ceftriaxone

Table 1: Antibacterial activity of 0.05 mg/ml ES of *Calliphore vomitoria* third larvae

Secretion/Excretion of uninfected bacteria	Growth zone (mm) of Inhibition			
	Gram- positive	Gram- Negatives		
	<i>S. aureus</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>K. pneumoniae</i>
Lower phase	18.5±0.5 a	12.2±0.8 bc	11.2± 1.3 b	10.3±1.5 c
Upper phase	17.2±1.2 a	12.8±0.3 c	10.8±0.3 cd	9.9±0.8 d
CRO (+ve)	9.2±0.8 a.	.3±1.0 c3	6.3±1.0 c	.3±1.0 c4

- Horizontal means ± SD with different letters are significantly different of P ≤ 0.05 (Duncan's test). CRO is Ceftriaxone antibiotic (standard drug).

Table 2: Growth inhibition of marker bacteria by antibacterial of 0.05 mg/ml ES of *Calliphore vomitoria* third larvae cannibalized bacteria infected larvae

E/S of infected larvae	Inhibition growth zone (mm) of			
	Gram- positive	Gram - Negatives		
	<i>S. aureus</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>K. pneumoniae</i>
Lower phase	28.3±0.6 b	21.8±1.0 c	33.2±1.0 a	17.3±1.5 c
Upper phase	25.0±0.5 a	17.2±0.8 b	23.2±0.8 ab	12.8±0.3 c
CRO (+ve)	9.2±0.8 a	3.3±1.0 c	.8±0.3 c	3.5±0.5 c

-Means in Horizontal view with different letters are significantly different letters are significantly different of P ≤ 0.05 (Duncan's test) -CRO is Ceftriaxone antibiotic (standard drug).

Table 3: Growth inhibition of marker bacteria by antibacterial of piperacillin resistant with ES of *Calliphore vomitoria* third larvae instar larvae at 0.1 and 0.05 mg/ml concentrations.

SE/Phase Conc. (mg/ml)		Inhibition growth zone (mm) of			
		Gram- positive	Gram- Negatives		
		<i>S. aureus</i>	<i>E. coli</i>	<i>P.aeruginosa</i>	<i>K. pneumoniae</i>
Clear phase	0.1	13.8±0.8 b	9.4±0.1c	16.8±0.5 a	14.0±0.8 ab
	0.05	10.0±0.5 a	5.5±1.5 bc	10.7±0.3 a	4.5±1.0 c
Dense phase	0.1	15.2±0.3 a	11.4±0.5 c	12.8±1.0 b	9.5±0.5 d
	0.05	9.2±0.8 a	6.9±0.5 b	8.8±0.8 a	4.2±1.5 c
CRO (+ve)		0.0±0.0	0.0±0.0	0.0±0.0	0.0±0.0

- Means in horizontal view with different letters are significantly different at P ≤ 0.05

Discussion

The results of this study are consistency with the hypothesis that the carrion insects were adapted with microbial stress and their immune system respond through anal excretions, oral excretions and internal secretions for inhibiting pathogenic agents: bacteria, fungi and yeasts. *Calliphora vomitoria* larvae

are obligate feeder on carrions. Our finding were supported that hypothesis within excretion/ secretion elution of the 3rd instar larvae by inert solvent (DMSO) and tested them against known sensitive and Piperacillin resistant bacteria. Positive control CRO antibiotic – standard drug - usage to evaluate antimicrobial action of the crude ES to build millstone for

next generation of alternative antibacterial molecules against today antibiotic resistant bacteria.

After ES fractionation, the two phases (dense and clear) were varied in an antibacterial activity against the Gram-positive *S. aureus* and Gram-negatives; *E. coli*, *P. aeruginosa* and *K. pneumoniae*. Table 1 shows that growth inhibition zones are in good effect according to [26] taxa, the clear phase was significantly more effective than dense phase, and more inhibited Gram-positive bacteria (*S. aureus*), than Gram-negatives (*E. coli*, *P. aeruginosa* and *K. Pneumoniae*). Clear and dense phases are more effective than the positive control (CRO) with valuable significance. In otherwise, inhibition zones of *P. aeruginosa* were 33.2 and 20.2 mm and for *K. Puenmoniae* 20.3 and 15.8 mm at clear and dense phases treatments, while those bacterial were not affected by CRO treatment. However dense and clear ES phases of stressed larvae which cannibalized infected larvae appeared high significant growth inhibitions (zone diameters), looking for the table 1 and 2 be saw the differences in growth inhibition zone values between the antimicrobial molecules so were from stressed (infected) or those without stress (uninfected) *C. vomitoria* larvae. The present finding were supported by

[10] and [1] were SE *L. sericata* and *C. vicina* inhibited growth of *S. aureus*, *P. aeruginosa* and *K. Pnuemoniae* and other microorganisms.

Today, resistant bacteria strains are demand inspecting for alternatives to solve problem of drug resistant. Therefore, figure 1 exhibits that ES of third larvae of *C. vomitoria* in either two applied phases (dense, clear) had meaningful efficacy against piperacillin resistant Gram positive and Gram negative marker bacterial strains in comparison with no effecting by CRO antibiotic. Our results were contest with [19], they found that induced immune activity of black soldier fly, *Hermtie illucens* larvae with contaminated needle, and isolated compounds from larvae extract caused antibacterial activity on sensitive and meticillin *S. aureus* and *P. aeruginose*. Otherwise, [17] were reported that insect body extracts have antibacterial effect on Multi drug Resistant bacteria. The present study was agreed with other articles were two AMPs identified from *Pichia pastoris* with high antimicrobial potency against Methicillin resistant *S. aureus* and other Gram positive bacteria [16], and feeding carrion and corpus maggots are produce a cocktail of protease enzymes and antimicrobial compounds called ES [25],




Fig 1: Relative antimicrobial activity of clear phase of larval SE fractionation of normal and infected fed larvae of *Calliphora vomitoria* at 0.05 mg/ml

Conclusion

The excretions / secretions (ES) of the carrion blue bottle fly, *Calliphora vomitoria* L. larvae were inhibited growth of the tested sensitive Gram – positive and Gram – negatives bacteria; *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella puenmoniae*. Also, ES inhibited growth of those previous bacterial species resistant piperacillin, and significantly that found better than standard drug (Ceftriaxone). The ES of the normal and others cannibalized infected and contaminated pathogenic larvae will encourage for more research in this topic.

Acknowledgment

The authors are grateful Mosul university authorities with anticipation to President of the university Prof. Dr. Kossay Alahmady for scientific support the first author. Also, thankful to Head of Biology Department; Assist. Prof. Dr. Mohmaad S. Faisal, for providing the research facilities in the Entomology laboratory/College of Education and Pure

Sciences, we thank the Bacteriologist Dr. Alaa Taha Younis for diligent help in practical part of this article.

References

1. Barends K, Gennard D, Dixon R. An assessment of the antimicrobial activity in larval excretion / with of four species of insects recorded in association with corpses, using *Lucilia sericata* Meigen as marker species. Bulletin of Entomological Research 2020;100(6):635-640.
2. Bauer AW, Kirby WM, Shemis C, Turck M. Antibiotic susceptibility testing by standardized single disk method. Am. J. Clin. Pathol 1966;45:493-496.
3. Bexfield A, Nijam Y, Thomas S, Ratcliffe NA. Detection and partial characterization of two antibacterial factors from the excretions / secretions of the medicinal maggot *Lucilia sericata* and their activity against methicillin – resistant *Staphylococcus aureus* (MRSA). Microbes Infect 2004;6:1297-1307.
4. Byrd JH, Castner JL. Forensic Entomology, the Utility of

- Arthropods in Legal Investigations 2nd ed. CRC Press 2010.
5. Cazander G, Pritchard DL, Nigam W, Nibbening PH. Multiple actions of *Lucillia sericata* larvae in hard – to – heal wound. *Bioassays* 35: 0000 – 0000, © 213 Wiley Periodicals, Inc 2013. Available from: DOI 10.1002/bies.201300071
 6. Chernysh SI, Gordja NA. The immune system of the blow fly (*Calliphora vicina*) as a source of medicinal drugs. *J. Evol. Biochem. Physiol* 2011;47(6):524-533.
 7. Elmorsy RH, Bream AS, Abdel – Samad RK. Antibacterial activity of *Chrysomya alpicops* maggot^s extracts (Diptera: Calliphoridae). *Egypt. Acad. J. Biolog. Sci* 2020;13(1):99-104.
 8. Golebiowski M, Cercowniak M, Urbanek A, Dawgul M, Kamysz W, Bogus MI *et al.* Antimicrobial activity of untypical lipid compounds in the cuticular and internal lipid of four fly species. *J. Appl. Microbiol* 2013;116:269-287.
 9. Hall C, Hadsworth NK, Howard DR, Jennings EM, Farrell LD, Magnuson TS *et al.* Inhibition of microorganisms on a carrion breeding resource: The antimicrobial peptide activity of Bunin beetle (Coleoptera: Silphidae) oral and anal secretions. *Environ. Entomol* 2011;40(3):669-625.
 10. Hassan MI, Amer MS, Hammad KM, Zidan MM. Antimicrobial activity for excretion and secretion of the green bottle fly larvae *Lucilia sericata* (Meigen) (Diptera: Calliphoridae). *J. Egypt. SOC. Parasitol* 2016;46(1):176-184.
 11. Horback WW, Bishop AA, Kreomer J, Scalzitty J, Scalzitti J, Shaffer JJ. Differences among antibacterial properties of carrion beetle secretions reflex phylogeny and Ecology. *J. Chem. Ecol* 2004;30(4):719-729.
 12. Jaklic D, Labanje A, Zupancic K, Smrake D, Gunde – Cimerman N. Selective antimicrobial activity of maggots against pathogenic bacteria. *J. Medical Microbiol* 2008;57(5):617-625.
 13. Januzanis B, Staczek S, Barbaras AZ, Badiul D, Gil JJ, Langer E *et al.* The effect of *Galleria mellonella* hemolymph polypeptides on human brain glioblastoma multi form cell line: A preliminary study. *Annals UMCS Biologia* 2012;67(2):52-62.
 14. Kojour MAH, Han YS, Jo YH. An overview of insect innate immunity. *Entomological research* 2020;50:282-291.
 15. Kurata S. Intra – and extracellular recognition of pathogens and activation of innate immunity. *Yakugaku Zasshi* 2006;126(12):1211-1218.
 16. Li Z, Mao R, Tang D, Hao Y, Chen H, Wang X *et al.* Antibacterial and immune modulatory activities of insect defenseins – DLP2 and DLP4 against multidrug – resistant *Staphylococcus aureus*. *Scientific Report* 2017;7:12124. DOI: 10.1038/s41598-017-10839-4
 17. Ma G, Wu L, Shao F, Zhang C, Wan H. Antibacterial activity of 11 insects extracts against multi drug resistant (MDR) strains of bacteria and fungus. *IOP Conf. Series: Earth Environmental Science* 2018;252(2019):022132. Available from: doi: 10.1088/1755-1315/2522/02213
 18. Mohamed NT. Activity of two serine protease and antibacterial studies on both excretory / secretory products and whole body homogenates of third instar larvae of *Chrysomya megacephala*. *Int. J. Adv. Rees* 2015;3(4):451-459
 19. Park S, Chang BS, Yoe SM. Detection of antimicrobial substances from larvae of the black soldier fly, *Heretia illucens* (Diptera: Stratiomyidae). *Entomological Research* 2014;44:58-64.
 20. Iamba K, Yoba S. Spatio-temporal dispersion patterns of *Bactrocera musae* Tryon (Diptera: Tephritidae: Dacinae) in Vudal agroecosystem, East New Britain. *International Journal of Entomology Research*. 2020;5(6):78-84.
 21. Poople A, Vogel H, Wiesner J, Vilicinskas A. Antimicrobial peptides expressed in medicinal maggots of the blow fly *Lucillia sericata* show combinatorial activity against bacteria, *Antimicrobial Agents and Chemotherapy* 2015;59:2508-2514.
 22. Sewify GH, Hammada HM, Alhadrami HA. *In vitro* evaluation of antimicrobial activity of alimentary canal excretions from the red palm weevil, *Rhynchophorus ferrugineus* Olivier larvae. *Biomed. Research International* 2017;2017(Article ID 856460):6. Available from: <https://doi.org/10.1555/2017/856460>
 23. Simmons SW. A bacterial principle in excretions of surgical maggots which destroys important etiological agents of pyogenic infections. *J. Bacteriol* 1935;30(3):253-267.
 24. Takov DI, Ostoich PV, Tchorbakov AI, Pilarska DK. Order Diptera as a model in the studies of insect immunity: a review. *Turkish Journal of Zoology* 2020;44:481-489.
 25. Thomas S, Andrews AM, Hay NP, Bourgoise S. The antimicrobial activity of maggot secretions: Results of a preliminary study. *J. Tissue Viability* 1999;14:97-101.
 26. Valachova I, Prochazka E, Bohova E, Novak P, Takac P, Majtan J. Antimicrobial properties of lucifersin in *Lucillia sericata* maggots after septic injury. *Asian Pac. J. Trop. Biomed* 2014;4(5):358-361.
 27. Van der Plas MJ, Jukema GN, Wai S, Bloemberg GV, Nibbering PH. Maggot excretions / secretions are differentially effective against biofilm of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *J. An. Microb. Chemotherapy* 2008;61:117-122.
 28. Vilcinskas A. Evolutionary plasticity of insect immunity. *J. Insect Physiolol* 2013;59:123-129.
 29. Wu Q, Patocha J, Kuka K. Insect antimicrobial peptides, a mini review. *Toxins* 2018;10:461.
 30. Yakovlev DY, Nesin AP, Simoenko NP, Gordya NA, Tulin DV, Kruglikova AA *et al.* Fat body and hemocytes contribution to the antimicrobial peptide synthesis in *Calliphora vicina* (Diptera: Calliphoridae) larvae. *In Vitro Cellular Developmental Biology – Animal* 2018;53:33-42.
 31. Pinilla YT, Patarroyo MA, Vlandia ML, Segura NA, Bello FJ. The effect of *Sarconesiopsis megellanica* larvae (Diptera: Calliphoridae) excretion and secretions on fibroblasts. *Acta Trop* 2015;142: 26-33.