



E-ISSN: 2320-7078

P-ISSN: 2349-6800

[www.entomoljournal.com](http://www.entomoljournal.com)

JEZS 2023; 11(4): 19-24

© 2023 JEZS

Received: 21-04-2023

Accepted: 20-05-2023

**Apoorva Indrapal Singh Kataria**  
Department of Ecosystem &  
Environment Management,  
Indian Institute of Forest  
Management, Bhopal, Madhya  
Pradesh, India

**Advait Edgaonkar**  
Department of Ecosystem &  
Environment Management,  
Indian Institute of Forest  
Management, Bhopal, Madhya  
Pradesh, India

## Forest insect pollination ecosystem services and climate change: A review

**Apoorva Indrapal Singh Kataria and Advait Edgaonkar**

DOI: <https://doi.org/10.22271/j.ento.2023.v11.i4a.9210>


### Abstract

Global warming has been predicted to cause big changes in terrestrial natural ecosystems and processes. Forests, being a part of terrestrial ecosystems, help in the regulation of biogeochemical processes and functions. Insect pollination is one function that is useful to the forest ecosystem as well as to the associated agroecosystems. Climate change is thought to be a principal factor that can influence the activities, distribution, and abundance of insect pollinators dwelling in the forest. Overall, elevated temperature level could affect the insect pollinators present in the forests. The present paper reviews the possible impact of climate change on insect pollination services provided by the forests, highlighting the importance of insect pollinators present in the forest, the dependence of linked agro-ecosystems on the forest, and various indicators of climate change with the species present in the forest. The paper finally concludes with some key recommendations for the conservation of forest insect pollination function with respect to climate change.

**Keywords:** Forest, pollinators, climate change, ecosystem services, agriculture

### 1. Introduction

Climate change has the potential to bring about pronounced shifts in insect behavior, their susceptibility to insecticides, and phenological synchrony with host plants (Pannure 2016) <sup>[48]</sup>. As per the IPCC report, temperature rise ranging from 1.1- 6.4 °C is estimated by the end of this century (Verghese *et al.* 2012) <sup>[61]</sup>. The forest forms an integral part of the biome encompassing a huge amount of biodiversity with a land cover of about 30% and an area of about 42 million km<sup>2</sup> in tropical, temperate, and boreal lands (Bonan 2008) <sup>[13]</sup>. Forests provide pollination as an ecosystem service of global importance providing significant economic and aesthetic benefits as well as cultural values to human society.



Source: Abrol, 2009 <sup>[2]</sup>

**Fig 1:** Contribution of insect groups pollinators to pollination

Insect pollinators show an astonishing taxonomic diversity in the forest ecosystem and are abundant in almost all environments across the globe. Hymenopterans with nearly 2,50,000 described species constitute the most important group of pollinators (Pannure 2016) <sup>[48]</sup>.


**Corresponding Author:**

**Apoorva Indrapal Singh Kataria**  
Department of Ecosystem &  
Environment Management,  
Indian Institute of Forest  
Management, Bhopal, Madhya  
Pradesh, India

Bees, flies, butterflies, moths, wasps, beetles, thrips, and some other insect orders encompass the majority of pollinating species (Fig.1).

Of the more than 100 crop species which provide 90% of the food supply for 146 countries, 71 are bee-pollinated (mainly

by wild bees) and several others are pollinated by wasps, flies, beetles, etc. (Pannure 2016) [48]. Native bee communities provide pollination services but the amount of benefit they provide and how this varies with land management practices is not well known.



**Fig 2:** Conceptual framework for understanding the flow of insect pollination service in the forest ecosystem

Over 85% of described flowering plant species are dependent, to some degree, on animal pollination, with mobile foraging insects accounting for the vast majority of this activity (Kremen *et al.* 2007) [39]. This results in increased fruit quality and quantity, seed production and fertility (Albrecht *et al.* 2012; Barrett 2003; Klein *et al.* 2007) [4, 7, 37].

This review attempts to understand the role and effect of forests in providing insect pollination ecosystem services and the possible factors responsible for bringing changes in the patterns of delivering insect pollination services influenced by climate change (Fig 2).

## 2. Indian context

In India, 116 species of Dipterans belonging to 16 families visiting 92 plant species have been reported. The most dominant group among the flower-visiting flies was the family Syrphidae, and Asteraceae was the most favored plant family (Mitra 2010, Sima *et al.* 2012) [45, 55].

The potential effects of climate change on pollinators have been reported by farmers practicing traditional beekeeping with *Apis cerana* in the Himalayas showing the effect of

temperature influencing the population sizes and swarming times of the insect pollinators (Tripathi 2011) [58]. In the Nainital district of Uttarakhand state the abundance of honeybees was significantly higher in an orchard having pollination services from apiary bees than in a control site where the pollination was dependent on available natural pollinators (Sharma 2012) [54].

The percent yield increase due to bee pollination for crops like mustard was found to be 43%, 32- 48% for sunflower, 17-19% for cotton, 93% for onion, and 44% for apple, etc. (Source: TNAU, Agrotech portal).

## 3. Effects of climate change on the forest insect pollination ecosystem services

Patterns of change in most pollinator assemblages suggest a causal connection between plants and their pollinators (Biesmeijer 2006) [12]. Studies on different climate change indicators are shown in Table. 1.

In the Kodagu district, India (Boreuxa *et al.* 2012) [15], the impact of forest fragments was observed. Here the bee visits and fruit set in rain-fed and irrigated coffee agro-forests were

seen affected by the forest and its size. Klein (2003) [36] found that coffee and cacao agro-forests in Sulawesi near rainforests had higher pollinator species diversity than in agro-forests situated further away, and as a consequence diversity was seen affecting the fruit-set more than abundance. Similarly in Brazil, farms near forest fragments had an increase in production up to 14.6% (De Marco & Coelho 2004) [25].

In tropical forests, the value of pollination services to agriculture is seen to increase coffee yields by 20% within 1 km of the forests, and pollination services performed by the insects improved coffee quality near the forest by reducing the frequency of "peaberries" by 27% (Ricketts 2004) [52].

#### 4. Indicators of climate change influencing insect pollinators

Insects are cold-blooded organisms, it is estimated that a 2 °C temperature shift could have a drastic effect on their behavior, abundance, life cycles, and reproductive capabilities, which may differ depending on pollinator size, age, and sex. (Yamamura and Kiritani 1998; Verghese *et al.* 2012) [62, 61]. Night-time temperatures may also have strong effects on the pollinator communities, affecting maturity and senescence, leading to differences in the time of nectar secretion and pistil receptivity. This can have an effect on the insect pollinators' visitation durations and crop pollination (Chabert *et al.* 2018;

Rohde and Pilliod 2021) [19, 53]. The metabolic rate of the insect pollinators may increase with the temperature (Willmer 1983) bringing changes in their flight activities. In honeybees, flight metabolic rate decreases with increasing temperature, and may be able to spend more time foraging when temperatures are rising. Large and dark-colored bees warm up and cool down faster than small and light-colored bees (Pereboom & Biesmeijer 2003) [49]. This results in differences in resource exploitation and might lead to shifts in pollinator assemblages with increasing temperatures (Taylor 1963) [57].

Relative humidity may affect an insect's visitation rates and nectar quality. The rate of equilibrium of nectar in the flower may also get influenced by humidity (Rohde and Pilliod 2021) [53].

The phenology of plants and their insect pollinators (bumblebees and whitefly) may be affected by the amount of light that is received. The reproductive success of bee-pollinated species is seen to be highly susceptible to seasonal changes in light in deciduous forests (Kudo 2008; Keeler *et al.* 2021) [40, 34]. There can be a possibility in some cases for rainfall being responsible for affecting the insect pollinators' abundance rates. Arnold *et al.* (2018) [5] found a positive relationship between the abundance of cocoa midge and rainfall.

**Table 1:** Literature showing the effect of climate change indicators on insect pollinators

Variables	Findings	References
Temperature	Effect on the abundance rate of the insect pollinators, developmental stages of plants and insects, and pollinators' efficiency. Insects are affected at spatial and temporal contexts.	Arroyo <i>et al.</i> (1994) [6]; Gordo & Jose (2006) [31]; Boyle-Makowski and Philogene (2012)
Humidity	Effect on insect visitation rates, nectar quality changes, nectar quality, etc.	Rohde and Pilliod (2021) [53]; Corbet <i>et al.</i> (1979) [22]; Corbet (1990) [21]
Light	Effect on the phenology of plants and insect pollinators.	Kudo (2008) [40]; Keeler <i>et al.</i> (2021) [34]
Rainfall	Effect on insect pollinators' abundance rates.	Arnold <i>et al.</i> (2018) [5]
Microclimate	Effect on timing of flower opening, and the exposure and receptivity of the style affecting insect pollinator's visitation rates.	Gols <i>et al.</i> (2021) [30]

Microclimate can affect small ectotherms like insects as they have limited ability to adjust under higher temperature ranges. Microclimate influences the timing of flower opening, extrusion and dehiscence, the survival and germination of pollen grains, and the exposure and receptivity of the style (Gols *et al.* 2021) [30]. The nectar quality of the flowers may be affected due to microclimate and insect visits. Microclimate inside the flower may influence the rate of equilibrium of nectar with the relative humidity of the air or the equilibrium concentration itself (Corbet *et al.* 1979) [22].

Factors affecting the phenology of insect pollinators are life cycle changes, alteration in the development rate, and availability of food resources (Bezerra *et al.* 2019 [11]; Tougeron 2019; Rohde and Pilliod 2021; Kendall *et al.* 2021) [53, 35]. Studies show a shift in the natural range of *Xylocopa* bees due to climate changes (Bartomeus *et al.* 2011; Bezerra *et al.* 2019) [11]. Climate change may alter the mutualism between plant flowering time and pollinator flight periods. There may be possibilities for phenological mismatches that contribute to the disruption of pollination of plants that flower later in the season. This could mainly influence specialist insect pollinators, though it may also reduce the generalist foragers.

In Eastern Massachusetts, New Hampshire, and South Africa the flower shape, temperature, light, and season appeared to be the most important variables influencing insect visitation

rates (Boyle-Makowski and Philogene 1985 [16]; Abrahamczyk *et al.* 2011; Bezerra *et al.* 2019) [11]. Reduced foraging activity and increased disease prevalence are due to changes in weather variables like temperature, humidity, and light quality/quantity (Rafferty and Ives 2011; Kendall *et al.* 2021) [35]. The reproductive success of certain pollinators is highly susceptible to seasonal fluctuations, and can result in decreased reproductive potential and adult mortality (Kudo *et al.* 2004; Kudo 2008; Bezerra *et al.* 2019 [11]; Kendall *et al.* 2021) [40, 41, 35].

#### 5. Loss in forest insect pollination service and effect on the agroecosystems

Klein *et al.* (2007) [37] showed that many crops do not directly rely on insect pollinators for the production of the part that is consumed, but pollinators are important for their propagation. Annual production of pollination services estimated worldwide is \$235 billion- \$577 billion (in 2015 US\$) which is 5-8% of market value directly linked with pollination services (Potts *et al.* 2016) [50]. As per the global surveys (Verghese *et al.* 2012) [61] on insect pollinators, the numbers of pests and diseases are more in horticultural crops than field crops, and they can affect the nutritional security of poor people under the influence of climate change.

In north-western Argentina, the diversity of flower visitors declined with decreasing forest fragment size, but the

frequency of flower visitors especially honey bee visits tended to increase (Aizen and Feinsinger 1994) [3]. Pollinator populations have declined across many regions due to habitat loss and degradation (Bezerra *et al.* 2019; Lee *et al.* 2021) [11]. Natural and restored fragments near Pondicherry, India, showed lower fruit sets due to the limitation of insect pollinators leading to low reproductive outputs (Nayak and Davidar 2010) [47].

Land-use intensification and disease have long driven pollinator losses (Bergholz *et al.* 2022) [10]. Vanbergen 2013 [60], Garibaldi *et al.* 2011) [28]. Due to climatic fluctuations, disease outbreak can occur in crops. The association between these different pressures is also likely are also likely to contribute to pollinator declines (Vanbergen 2013) [60].

Studies on biological invasions showed that invasions affected insect pollinators by decreasing visitation rates and the fruit set percentage (Kaiser-Bunbery 2011; Montero-Castano and Vila 2012) [33, 46]. Studies suggest that plant invasion may attract fewer floral visitors (Montero-Castano & Vila 2012) [46] affecting the pollination activity (Brown *et al.* 2002) [18].

Frequent use of pesticides on crops may lead to reduced foraging activity of insect pollinators (Kendall *et al.* 2021) [35]. Surveys on wild bees and butterflies show a lower species richness in the places where the use of pesticides is more (Brittain *et al.* 2010) [17]. Systemic pesticides i.e., neonicotinoids have the capacity to penetrate through plant tissue resulting in accumulation in the plant nectar and pollen. This can result in sublethal effects on insect pollinators' activity and behavior (Cresswell 2011) [23]. It could affect the brain functioning, growth, and foraging capacity of honey bees (Vanbergen 2013, Gill *et al.* 2012) [60].

Droughts can lead to modifications in floral signals and rewards which could impact the reproductive success of both plants and pollinators negatively by decreasing pollen viability, pollen tube growth, and pollen deposition effectiveness. (Fang *et al.* 2010; Vaudo *et al.* 2015; Solis-Montero and Vallejo-Marin 2017; Arnold *et al.* 2018; Descamps *et al.* 2021) [27, 46, 5, 26].

N-fixing bacteria allow the plants to produce higher-quality rewards by increasing phosphorous and micronutrients, increasing pollinator visitation. A reduction in the strength of the interaction between legumes and rhizobia can affect plant nitrogen negatively (Keeler *et al.* 2021) [34]. This can affect pollinator feeding preference due to reduced flower number, nectar, and pollen quality. This could in turn have an effect on bee fitness. Potassium is also an important micronutrient for bee reproductive success (Praz *et al.* 2008; Davis *et al.* 2019; Bonoan *et al.* 2018; McAulay *et al.* 2021) [24, 14].

Fires regimes are seen to be impacting the functions of insect pollinator communities. Insect pollinator responses to fire are tied to multiple factors like fire history, plant responses, and changes in species interactions. Due to spatiotemporal and historical heterogeneity of fires and interaction of global drivers of insect pollinators change, prediction of community-level resilience of insect pollinators is often challenging (Koltz *et al.* 2018) [38].

The artificial night light is seen as a reason for reduced population sizes and changes in the community composition of insect pollinators. It can lead to disruption of visual cues and affect pollinators' behavior and populations. It could also result in affecting the plant-pollinator interaction patterns (Grubisic and Grunsven 2021; Macgregor *et al.* 2017; Van Grunsven *et al.* 2020) [32, 43, 59].

## 6. Conclusions

Climate change may adversely affect ecosystem functions

therefore managing forests' ecosystems can prove beneficial in mitigating the adverse effects of climate change. The effects of climate change on insect pollinator species may have positive as well as negative consequences.

Out of all the climatic variables affecting insect pollinators, the temperature is found to be the most prominent factor influencing the forest ecosystem. Rise and fall in temperature levels more than the normal range could create an enormous impact on insect pollinator diversity and abundances, their phenology, and their distribution patterns. Because of fluctuation in the temperature ranges every year, there may be chances of reduced floral appearances, delayed flowering and fruiting, disease outbreaks, etc. It may also cause a shift in the pattern of insect pollinators occurrences, morphology, and their habitat range, etc. studies show that it is due to climate change that an insect pollinator species dwelling in a particular ecosystem may alter the fundamental niche habitat. It may enable survival of that species but the plants dependent on these insect pollinators may get affected adversely.

Globally and in India much research needs to be performed on insect pollinators and their effect on forests and the agroecosystem. While the attributes affecting climate change are difficult to analyze, there are still some parameters and data available, explaining the interrelationships between the climate variables affecting insect pollination service. However, the scale of dependence of plants and tree species present in the forest ecosystem on the insect pollinators is not known.

Despite knowledge of importance of insect pollinators and their role in providing the service to the agroecosystem for crop production, there is still limited data on species diversity, abundance, and community composition of insect pollinators. There is a need to carry out research on the forest floral species-specific insect pollinator. One of the major problems is the availability of baseline data. Therefore, in order to reduce the research gap, it is important that more attention be given to baseline data generation on the insect pollinators, their diversity, and abundances. This ground data gathered over years could be utilized in understanding the plant species-specific insect pollinators and comparing it with climatic events using some climatic models. Thus the impact of climate change on both plants as well as insect pollinators could be predicted.

## 7. Acknowledgments

The author would like to thank the Madhya Pradesh government and EPCO, Bhopal for funding the research under the Chief Minister's Scholarship for Ph.D. on Climate Change. This study is part of the same research work.

## 8. References

1. Abrahamczyk S, Kluge J, Gareca Y, Reichle S, Kessler M. The influence of climatic seasonality on the diversity of different tropical pollinator groups. *PLOS One*. 2011;6(11):e27115.
2. Abrol DP. Plant-pollinator interactions in the context of climate change-an endangered mutualism. *Journal of Palynology*. 2009;45:1-25.
3. Aizen MA, Feinsinger P. Habitat fragmentation, native insect pollinators and feral honeybees in Argentine' Chaco Serrano'. *Ecological Applications*. 1994;4(2):378-392.
4. Albrecht M, Schmid B, Hautier Y, Mueller CB. Diverse pollinator communities enhance plant reproductive

- success. Proc. R. Soc. B Biol. Sci. 2012;279:4845–4852.
5. Arnold SEJ, Bridgemohan P, Perry GB, Spinelli GR, Pierre B, Murray F. The significance of climate in the pollinator dynamics of a tropical agroforestry system. *Agriculture, ecosystems & environment*. 2018;254:1-9.
  6. Arroyo MTK, Armesto JJ, Primack RB. Community studies in pollination ecology in the high temperate Andes of central Chile II. Effect of temperature on visitation rates and pollination possibilities. *Plant systematics and evolution*. 1985;149(3-4):187-203.
  7. Barrett SCH. Mating strategies in flowering plants: the outcrossing-selfing paradigm and beyond. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 2003;358:991–1004.
  8. Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R. Climate-associated phenological advances in bee pollinators and bee-pollinated plants. *Proceedings of the National Academy of Sciences*. 2011;108(51):20645-20649.
  9. Gurung S, Mandal RA, Mathema AB. Valuation of ecosystem services provided by Jal Binayak community forest, Nepal. *Int. J Geogr Geol. Environ* 2021;3(2):25-34.
  10. Bergholz K, Sittel LP, Ristow M, Jeltsch F, Weiss L. Pollinator guilds respond contrastingly at different scales to landscape parameters of land-use intensity. *Ecology and evolution*. 2022;12:e8708.
  11. Bezerra ADM, Pacheco Filho AJ, Bomfim IG, Smagghe G, Freitas BM. Agricultural area losses and pollinator mismatch due to climate changes endanger passion fruit production in the Neotropics. *Agricultural systems*. 2019;169:49-57.
  12. Biesmeijer JC, Roberts SP, Reemer M, Ohlemuller R, Edward M, Peeters T. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*. 2006;313(5785):351-354.
  13. Bonan GB. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science*. 2008;320(5882):1444-1449.
  14. Bonoan RE, O'Connor LD, Starks PT. Seasonality of honey bee (*Apis mellifera*) micronutrient supplementation and environmental limitation, *J Insect Physiol*. 2018;107:23–28.
  15. Boreux V, Krishnan S, Cheppudira KG, Ghazoul J. Impact of forest fragments on bee visits and fruit set in rain-fed and irrigated coffee agro-forests. *Agriculture, ecosystems and environment*. 2013;172:42-48.
  16. Boyle-Makowski RMD, Philogene BJR. Pollinator activity and abiotic factors in an apple orchard. *The Canadian Entomologist*. 1985;117(12):1509-1521.
  17. Brittain CA, Vighi M, Bommarco R, Settele J, Potts SG. Impacts of a pesticide on pollinator species richness at different spatial scales. *Basic and Applied Ecology*. 2010;11:106-115.
  18. Brown BJ, Mitchell RJ, Graham SA. Competition for pollination between an invasive species (Purple loosestrife) and a native congener. *Ecology*. 2002;83:2328-2336.
  19. Chabert S, Lemoine T, Cagnato MR, Morison N, Vaissiere BE. Flower age expressed in thermal time: is nectar secretion synchronous with pistil receptivity in oilseed rape (*Brassica napus* L.)? *Environmental and Experimental Botany*. 2018;155:628-640.
  20. Corbet SA. Conserving compartments in pollination webs. *Conservation biology*. 2000;14:1229-1231.
  21. Corbet SA. Pollination and the weather. *Israel Journal of Plant Sciences*. 1990;39(1-2):13-30.
  22. Corbet SA, Unwin DM, Prys-Jones OE. Humidity, nectar and insect visits to flowers, with special reference to *Crataegus*, *Tilia* and *Echium*. *Ecological Entomology*. 1979;4(1):9-22.
  23. Cresswell JE. A meta-analysis of experiments testing the effects of a neonicotinoid insecticide (imidacloprid) on honey bees. *Ecotoxicology*. 2011;20:149-157.
  24. Davis JK, Aguirre LA, Barber NA, Stevenson PC, Adler LS. From plant fungi to bee parasites: Mycorrhizae and soil nutrients shape floral chemistry and bee pathogens. *Ecology*. 2019;100:e02801.
  25. De Marco P, Coelho FM. Services performed by the ecosystem: Forest remnants influence agricultural cultures' pollination and production. *Biodiversity & Conservation*. 2004;13(7):1245-1255.
  26. Descamps C, Maree S, Hugon S, Quinet M, Jacquemart AL. Species-specific responses to combined water stress and increasing temperatures in two bee-pollinated congeners (*Echium*, Boraginaceae). *Ecology and Evolution*. 2020;10:6549-6561.
  27. Fang X, Turner NC, Yan G, Li F, Siddique KH. Flower numbers, pod production, pollen viability, and pistil function are reduced and flower and pod abortion increased in chickpea (*Cicer arietinum* L.) under terminal drought. *Journal of experimental botany*. 2010;61:335-345.
  28. Garibaldi LA, Steffan-Dewenter I, Kremen C, Morales JM, Bommarco R, Cunningham SA. Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology letters*. 2011;14:1062-1072.
  29. Gill RJ, Ramos-Rodriguez O, Raine NE. Combined pesticide exposure severely affects individual and colony-level traits in bees. *Nature*. 2012;491:105-108.
  30. Gols R, Ojeda-Prieto LM, Li K, Van Der Putten WH, Harvey JA. Within-patch and edge microclimates vary over a growing season and are amplified during a heatwave: Consequences for ectothermic insects. *Journal of Thermal Biology*. 2021;99:103006.
  31. Gordo O, Sanz, JJ. Temporal trends in phenology of the honey bee *Apis mellifera* (L.) and the small white *Pieris rapae* (L.) in the Iberian Peninsula (1952–2004). *Ecological Entomology*. 2006;31(3):261-268.
  32. Grubisic M, Van Grunsven RH. Artificial light at night disrupts species interactions and changes insect communities. *Current opinion in insect science*. 2021;47:136-141.  
[http://www.agritech.tnau.ac.in/about\\_us/abt\\_us.html](http://www.agritech.tnau.ac.in/about_us/abt_us.html)
  33. Kaiser-Bunbury CN, Valentin T, Mougou J, Matatiken D, Ghazoul J. The tolerance of island plant-pollinator networks to alien plants. *Journal of Ecology*. 2011;99(1):202-213.
  34. Keeler AM, Rose-Person A, Rafferty NE. From the ground up: Building predictions for how climate change will affect belowground mutualisms, floral traits, and bee behavior. *Climate Change Ecology*. 2021;1:100013.
  35. Kendall LK, Evans LJ, Gee M, Smith TJ, Gagic V, Lobaton JD, Rader R. The effect of protective covers on pollinator health and pollination service delivery. *Agriculture, Ecosystems & Environment*. 2021;319:107556.

36. Klein AM, Steffan-Dewenter I, Tscharntke T. Pollination of *Coffea canephora* in relation to local and regional agroforestry management. *Journal of Applied Ecology*. 2003;40(5):837-845.
37. Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society of London B: Biological Sciences*. 2007;274:303-313.
38. Koltz AM, Burkle LA, Pressler Y, Dell JE, Vidal MC, Richards LA. Global change and the importance of fire for the ecology and evolution of insects. *Current opinion in insect science*. 2018;29:110-116.
39. Kremen C, Williams NM, Aizen MA, Gemmill-Herren B, Lebuhn G, Minckley R, *et al*. Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecol. Lett*. 2007;10:299–314.
40. Kudo G, Ida TY, Tani T. Linkages between phenology, pollination, photosynthesis, and reproduction in deciduous forest understory plants. *Ecology*. 2008;89(2):321-331.
41. Kudo G, Nishikawa Y, Kasagi T, Kosuge S. Does seed production of spring ephemerals decrease when spring comes early? *Ecological Research*. 2004;19(2):255-259.
42. Lee MR, Mc Neil Jr. DJ, Mathis CL, Grozinger CM, Larkin JL. Microhabitats created by log landings support abundant flowers and insect pollinators within regenerating mixed-oak stands in the Central Appalachian Mountains. *Forest Ecology and Management*. 2021;497:119472.
43. Macgregor CJ, Evans DM, Fox R, Pocock MJ. The dark side of street lighting: Impacts on moths and evidence for the disruption of nocturnal pollen transport. *Glob Change Biology*. 2017;23:697-707.
44. Mc Aulay MK, Killingsworth SZ, Forrest JRK. Understanding pollen specialization in mason bees: a case study of six species. *Oecologia*, 195, 559–574.
45. Mitra. Diversity of flower-visiting flies (Insecta: diptera) in India and their role in pollination. *Records of the Zoological Survey of India*. 2021;110(2):95-107.
46. Montero-Castano A, Vila M. Impact of landscape alteration and invasions on pollinators: A meta-analysis. *Journal of Ecology*. 2012;100(4):884-893.
47. Nayak KG, Davidar P. Pollinator limitation and the effect of breeding systems on plant reproduction in forest fragments. *Acta Oecologica*. 2010;36(2):191-196.
48. Pannure A. Bee pollinators decline Perspectives from India. *International research journal of natural and applied sciences*. 2016;3(5):2349-4077.
49. Pereboom JJ, M Biesmeijer JC. Thermal constraints for stingless bee foragers: the importance of body size and coloration. *Oecologia*. 2003;137(1):42-50.
50. Potts SG, Imperatriz-Fonseca VL, Ngo HT, Biesmeijer JC, Breeze TD, Dicks LV. Summary for policymakers of the assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production. 2016; Report No.: hal-01946814.
51. Rafferty NE, Ives AR. Effects of experimental shifts in flowering phenology on plant–pollinator interactions. *Ecology Letters*. 2011;14(1):69-74.
52. Ricketts TH, Daily GC, Ehrlich PR, Michener CD. Economic value of tropical forest to coffee production. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;101(34):12579-12582.
53. Rohde AT, Pilliod DS. Spatiotemporal dynamics of insect pollinator communities in sagebrush steppe associated with weather and vegetation. *Global Ecology and Conservation*. 2021;29:e01691.
54. Sharma HK, Partap U, Bisht K, Tewari P, Phartiyal P. Impact of honeybee pollination in enhancing the apple production in Nainttal district of Uttarakhand. *Adv. Pollen Spore Res*. 2012;30:99-102.
55. Sima DB, Srivastava M. Floral visitors of different crops as recorded from an agro-ecosystem near Jhunjhunu, Rajasthan (India). *Int. J Sci. Res*. 2014;3(9):1732-1738.
56. Solis-Montero L, Vallejo-Marin M. Does the morphological fit between flowers and pollinators affect pollen deposition? An experimental test in a buzz-pollinated species with another dimorphism. *Ecology and Evolution*. 2017;7:2706-2715.
57. Taylor LR. Analysis of the effect of temperature on insects in flight. *The Journal of Animal Ecology*. 1963;99-117.
58. Tripathi H. Role of beekeeping with indigenous bee- *Apis cerana* in crop production. *Beekeeping & agricultural productivity. Under the Mango Tree*. 2011, 36.
59. Van Grunsven RHA, Van Deijk JR, Donners M, Berendse F, Visser ME, Veenendaal E, Spoelstra K. Experimental light at night has a negative long-term impact on macro-moth populations. *Current Biology*. 2020;30:694- 695.
60. Vanbergen AJ. Threats to an ecosystem service: pressures on pollinators. *Frontiers in Ecology and the Environment*. 2013;11(5):251-259.
61. Verghese A, Reddy PR, Chandrashekar KN, Roopa HK. Climate change scenario in horticultural crops with reference to insect pests, biotypes, pollinators, and natural enemies. 2012.
62. Yamamura K, Kiritani KA. Simple method to estimate the potential increase in the number of generations under global warming in temperate zones. *Applied Entomology and Zoology*. 1998;33:289-298.
63. Ghosh S. Climate change and earthworms: A global perspective. *International Journal of Entomological Research*. 2021;6(4):167-71.
64. Batra SW. Biology of *Apis laboriosa* Smith, a pollinator of apples at high altitude in the great Himalaya range of Garhwal, India, (Hymenoptera: Apidae). *Journal of the Kansas Entomological Society*. 1996, 177-181.