Journal of Entomology and Zoology Studies

Available online at www.entomoljournal.com

Journal of Entomology and Zoology Studies

E-ISSN: 2320-7078
P-ISSN: 2349-6800
www.entomoljournal.com
JEZS 2025; 13(4): 33-36

Received: 22-04-2025
Accepted: 26-05-2025

Udaya Sree M
Department of Entomology, College of Horticulture, Dr. YSRHU, Andhra Pradesh, India

Nandini B
Department of Entomology, College of Agriculture, Rajendranagar, PJTAU, Hyderabad, Telangana, India

Corresponding Author:
Udaya Sree M
Department of Entomology, College of Horticulture, Dr. YSRHU, Andhra Prades, India

Semiochemicals: Role in IPM

Udaya Sree M and Nandini B

DOI: https://www.doi.org/10.22271/j.ento.2025.v13.i4a.9537

Abstract
Insect pests communicate and interact with their surroundings through a diverse array of chemical signals. Semiochemicals particularly pheromones have emerged as critical tools in environmentally sound Integrated Pest Management (IPM). Plants and insects have coevolved for over 350 million years, relying on intricate chemical signaling-including insect semiochemicals (e.g., pheromones) and plant defense compounds-to communicate, defend, and thrive. Insects leverage semiochemicals for behaviors like mating, host-finding, and predator avoidance, enabling precise, species-specific pest management strategies such as mating disruption, mass trapping, and attract-and-kill. Meanwhile, plants utilize constitutive and induced defenses-ranging from structural traits and stored toxins to herbivore-induced volatiles (HIPVs)-to deter herbivores and recruit natural enemies. Constitutive defenses include barriers like thorns and trichomes, and toxin storage strategies exemplified by the glucosinolate-myrosinase "mustard oil bomb," which rapidly generates toxic isothiocyanates upon tissue damage. Induced mechanisms, triggered by insect attack, activate internal signaling cascades that lead to secondary metabolite production, proteinase inhibitors, and volatile emissions-both at the local site and systemically-to deter attackers and alert neighboring plants. These volatiles, derived from pathways like terpenoids and fatty acids, have been shown in multiple species (e.g., maize, cotton, tobacco) to modulate herbivore behavior and attract parasitoids and predators. Modern IPM integrates these biochemical insights with advanced delivery technologies-such as controlled-release dispensers (e.g., SPLAT), smart traps, and genetically engineered volatile production in plants-to optimize pest control while preserving beneficial organisms and reducing pesticide reliance. However, realizing the full potential of semiochemical-based strategies requires enhanced formulation, deployment methods, ecological understanding, and interdisciplinary innovation spanning chemistry, molecular biology, and field ecology. As global agriculture seeks sustainable alternatives to synthetic pesticides, semiochemicals offer smart, precise, and environmentally friendly tools for future pest management.

Keywords: Semiochemicals, pheromones, IPM, mating disruption, mass trapping, push-pull strategy, chemical ecology

Introduction
Chemical signaling explains a wide range of insect behaviors, from mate and food finding to predator defense. Because insects are small and rely less on visual or auditory cues, they depend heavily on semiochemicals like pheromones for communication. Discoveries in these signals have led to sustainable, species-specific pest control strategies (e.g., mating disruption, mass trapping) that form the bedrock of modern chemical ecology and integrated pest management. Plants and insects have coexisted for over 350 million years, evolving a complex web of interactions. Some are mutually beneficial-such as pollination and seed dispersal, while others, like herbivory, are detrimental to plants. To defend themselves, plants have evolved diverse chemical defenses at all trophic levels, producing toxic or unpalatable compounds that significantly inhibit insect feeding.

These defenses begin with precise detection: plants recognize insect-derived molecules, triggering internal signaling cascades that activate responses ranging from physical barriers to the synthesis of secondary metabolites-such as phenylpropanoids, alkaloids, terpenoids, and fatty acid derivatives. Many compounds are volatile, allowing plants to engage in tritrophic communication-attracting predators or parasitoids of herbivores and facilitating inter-plant signaling. Building on this natural chemical ecology, researchers are now engineering plants or microbes to produce insect pheromones or herbivore-induced plant volatiles (HIPVs), enabling innovations like push-pull cropping systems, mating disruption, and attract-and-kill strategies.

These tactics reduce reliance on broad-spectrum pesticides and enhance natural pest resistance in crops. Through millions of years of coevolution, plants and insects have developed intricate chemical communications-plants using chemical detection and defense, and insects responding via semiochemicals. By harnessing these natural signaling systems (e.g., pheromones, HIPVs), modern agriculture is embracing environmentally friendly, targeted pest control-pioneering a sustainable future in crop protection.

Defence strategies: As primary producers, plants are consumed by a diverse array of herbivores-including nematodes, arthropods, molluscs, and mammals. They have evolved a broad spectrum of physicochemical defenses to protect themselves. These defenses fall into two categories: constitutive (always present) and induced (activated after attack). Constitutive defenses encompass structural and morphological traits, as well as innate chemical compounds that provide constant protection. Induced defenses activate upon herbivore or pathogen assault, and can be further divided into: Direct defenses, which impair the herbivore's survival or performance-such as accumulation of toxins or deterrents, cell wall reinforcement, and hypersensitive responses. Indirect defenses, which enlist a third trophic level by modifying the behaviors of predators or parasitoids to defend the plant. These may include the production of extrafloral nectar, food bodies, or domatia to provide food and shelter for carnivores. Both direct and indirect defenses can be either constitutive or induced, allowing plants to balance ongoing protection with resource-efficient responses to environmental threats.

Constitutive physical defenses: Constitutive physical defenses in plants range from visible deterrents to microscopic barriers: thorns and prickles ward off larger herbivores, while thick cell walls, robust cuticles, high lignin content, and resin accumulation impede pathogens and small herbivores. Trichomes-plant hairs-serve as mechanical defenses, and certain glandular types secrete antimicrobial substances, feeding deterrents, or toxins. Additionally, plants produce digestibility-reducing compounds such as tannins, resins, proteinase inhibitors, and silica; these anti-nutritional chemicals impair herbivore digestion, slowing growth, reducing fecundity, and increasing disease susceptibility. The effectiveness of toxic compounds is often linked to their storage and deployment: many species sequester toxins in specialized structures like resin ducts or glandular trichomes, which release defensive chemicals when physically damaged. To prevent self-harm, plants typically avoid storing active toxins. Instead, they accumulate inactive precursors-such as glycosides-separated from their activating enzymes. Upon tissue damage, enzymes like myrosinase mix with the precursors (e.g., glucosinolates), triggering the release of toxic compounds such as isothiocyanates or nitriles, in a well-documented defense mechanism known as the "mustard oil bomb." Individual toxins further exhibit unique modes of action: saponins disrupt cellular membranes, cyanogenic glycosides release hydrogen cyanide that inhibits respiration, cardenolides target the Na^+/K^+-ATPase pump, and certain polyacetylenes, like cicutoxin, interfere with neuronal potassium channels. Together, these layered structural and chemical defenses form a robust and comprehensive protective system against a wide range of herbivores and pathogens.

Induced defenses: In contrast to constitutive defenses, **induced defenses** require early detection of an insect attack and activation of alarm signaling to mount a fast and effective response. These plant strategies-which may be constitutive, inducible, or a combination-frequently produce systemic effects, extending beyond the initial damage site to the rest of the plant. Such responses involve the synthesis and accumulation of defensive secondary metabolites that deter insect feeding and disrupt development. Additionally, proteins like proteinase inhibitors (PIs), polyphenol oxidases (PPOs), and threonine deaminase (TD) inhibit insect digestive enzymes or reduce the nutritional quality of plant tissues, acting as direct inducible defenses. When attacked by herbivores, many plants emit volatile organic compounds (VOCs) that serve multiple functions: guiding herbivores to their hosts, recruiting predators or parasitoids (an indirect defense involving a third trophic level), and signaling other parts of the same or neighboring plants to bolster their defenses. These volatiles-comprising terpenoids (mono-, di-, sesqui-, and homoterpenoids), C6 fatty acid derivatives (e.g., (3Z)-hex-3-enal, (3Z)-hex-3-enyl acetate), aromatic molecules like methyl salicylate and indole, plus various alkanes, alkenes, and alcohols-form a dynamic "volatilome" shaped by complex biosynthetic pathways responsive to environmental cues.

Numerous plants, such as maize (*Zea mays*), cotton (*Gossypium hirsutum*), lima bean (*Phaseolus lunatus*), *Lotus japonicus*, tobacco (*Nicotiana attenuata*), and *Medicago truncatula*, have been shown to emit herbivore-induced volatiles (HIVs). These compounds influence plant-insect community interactions by modifying herbivore behavior and attracting natural enemies.

Recent research has revealed novel roles for these HIVs beyond long-range attraction. For example, after visiting aphid-infested cucumber (*Cucumis sativus*) plants, parasitoids (*Lysiphlebus testaceipes*) spend less time on and attack fewer aphids in subsequent patches-not due to the presence of aphids, but because of plant-emitted volatiles, indicating that parasitoids adapt their foraging based on previously encountered cues. As a result, plants emitting stronger volatile signals may attract and retain more parasitoids, creating positive selection pressure for high emission rates. Volatile emissions often correlate with herbivore density, helping predators and parasitoids preferentially target plants under heavier attack. By identifying electrophysiologically-and behaviorally-active compounds through techniques like gas chromatography-electroantennography, scientists can pinpoint specific molecules (often from the isoprenoid and lipoxygenase pathways, such as (E)-ocimene and (-)-germacrene D) that repel herbivores or attract their enemies.

Below ground, volatiles are equally important. Maize roots damaged by the pest *Diabrotica virgifera virgifera* emit (E)-β-caryophyllene, which attracts entomopathogenic nematodes to attack the insect, a defense conspicuously absent in varieties lacking the compound. Furthermore, herbivory or jasmonic acid treatment in lima beans triggers both volatile emission and the secretion of extrafloral nectar, which attracts protective ants. This defense strategy reduces leaf damage and improves plant fitness, benefits that also spread to neighboring plants through induced leaflet signaling.

Together, these findings illustrate how volatile-mediated signals-above and below ground-shape interactions within plant-insect communities, offering promising avenues for developing pest-resistant crops through ecological

engineering of VOC emissions.

Functional Categories of Semiochemicals

Cuticular Hydrocarbons (CHCs)

Insect cuticles bear hydrocarbon layers that originally evolved to prevent desiccation. Over time, these compounds began serving additional roles-especially in sexual signaling and social identification. CHCs differ by species, sex, and physiological status, making them reliable tools for pest monitoring and behavior manipulation.

Multifunctional Defensive Secretions

Certain insects use chemical defenses not only to deter predators but also to communicate. For instance, formic acid in ants and (E,E)-α-farnesene in termites act as both repellents and alarm pheromones. In some beetles and parasitoids, defensive substances also attract mates, highlighting their dual ecological roles.

Hormone-Derived Compounds

Though less common, some pheromones originate from hormone pathways. The burying beetle Nicrophorus vespilloides emits methyl geranate-a compound linked to the juvenile hormone pathway-which temporarily suppresses fertility and fosters cooperative parenting.

Plant-Origin Semiochemicals

Some moths and butterflies ingest and convert specific plant alkaloids (e.g., pyrrolizidine alkaloids) into sex pheromones. These compounds not only facilitate mating but also enhance offspring defense, indicating a strong evolutionary link between diet and reproductive signaling.

Social Structure and Colony Dynamics

In eusocial species like ants and bees, chemical signals govern caste roles and colony behavior. Queen pheromones suppress worker reproduction, trail pheromones aid foraging, and alarm signals synchronize defense-all crucial for maintaining colony stability.

Tri-Trophic Chemical Signaling

Plant-emitted volatiles triggered by herbivore attack (HIPVs) can attract natural enemies like parasitoids. Leveraging such tri-trophic cues in IPM can significantly improve conservation biological control and reduce pesticide dependence.

Applications in Pest Management

Pest Monitoring and Surveillance

Pheromone-baited traps are vital for detecting pest presence, population trends, and seasonal activity. These tools guide timely interventions, reducing unnecessary pesticide applications.

Mating Disruption

Saturating the environment with synthetic sex pheromones misleads male insects, preventing them from locating mates. Though mating isn't completely blocked, the resultant delay lowers reproductive success and pest proliferation.

Attract-and-Kill Techniques

By combining attractants with lethal agents, pests are drawn to specific zones where they are exterminated. This technique minimizes crop exposure to insecticides while effectively curbing pest populations.

Mass Trapping

Deploying large numbers of pheromone traps helps reduce pest densities, particularly for species with limited reproductive output. It is widely used for managing pests like the red palm weevil.

Push-Pull Strategy and Repellents

This method uses repellent compounds to drive pests away from crops ("push") and attractants to lure them into trap areas ("pull"). Understanding pest behavior is crucial for optimizing this approach.

Enhancing Biological Control

Semiochemicals like kairomones can attract beneficial parasitoids or predators to pest hotspots, enhancing the success of biological control. Baited refuges can also support natural enemy populations in agroecosystems.

Conclusion

Semiochemicals play a pivotal role in modern, sustainable pest management by precisely disrupting pest communication and bolstering natural enemies-reducing dependency on conventional pesticides. These compounds are employed in diverse strategies-monitoring (to inform control decisions), mating disruption, mass trapping, and attract-and-kill-each meticulously designed to manipulate insect behavior. Monitoring using pheromone traps guides timely interventions. Mating disruption, via synthetic pheromones, prevents pest reproduction, while attract-and-kill techniques lure pests to treated surfaces for elimination. These strategies are part of integrated pest management (IPM) frameworks that merge semiochemicals with biological control, cultural practices, crop resistance, and precision agriculture tools to achieve targeted, eco-friendly pest suppression.

Although research has validated the efficacy of semiochemical approaches, practical adoption remains limited. Future efforts must focus on refining formulations and delivery systems-such as controlled-release dispensers like SPLAT and puffers (MSTRS)-to improve field performance and durability. Integrating semiochemicals with precision farming technologies (e.g., remote sensing, smart traps) can optimize timing and localization of deployment. Developing behavior-modifying agents calls for deep understanding of biochemical sources, and molecular, physiological, and ecological interactions between plants, pests, and natural enemies. Cutting-edge methods-such as synthesizing pheromones in plants, auto-dissemination of entomopathogens, and "plant vaccination"-show promise for highly targeted, species-specific chemical signaling.

Advancing semiochemical-based IPM demands not only technical innovation but also solid ecological understanding to ensure efficacy, avoid non-target effects, and integrate seamlessly into cropping systems. As challenges with synthetic pesticides escalate-health risks, resistance, and environmental impact-behavior-modifier tools grounded in ecological and molecular science stand out as scalable, smart, and sustainable alternatives.

References

1. Aljbory Z, Chen M-S. Indirect plant defense against insect herbivores: a review. Insect Sci. 2018;25:2-23. Available from: https://doi.org/10.1111/1744-7917.12436
2. Chung SH, Rosa C, Scully ED, Peiffer M, Tooker JF, Hoover K, *et al*. Herbivore exploits orally secreted

bacteria to suppress plant defenses. Proc Natl Acad Sci. 2013;110:15728-1533. Available from: https://doi.org/10.1073/pnas.1308867110
3. Holopainen JK, Blande JD. Where do herbivore-induced plant volatiles go? Front Plant Sci. 2013;4:1-13. Available from: https://doi.org/10.3389/fpls.2013.00185
4. Compant S, Samad A, Faist H, Sessitsch A. A review on the plant microbiome: ecology, functions, and emerging trends in microbial application. J Adv Res. 2019;19:29-37. Available from: https://doi.org/10.1016/j.jare.2019.03.004
5. Mello MO, Silva-Filho MC. Plant-insect interactions: an evolutionary arms race between two distinct defense mechanisms. Bras J Plant Physiol. 2002;14:71-81. Available from: https://doi.org/10.1590/S1677-04202002000200001
6. Midega CAO, Pittchar JO, Pickett JA, Hailu GW, Khan ZR. A climate-adapted push-pull system efectively controls fall armyworm, *Spodoptera frugiperda* (J E Smith), in maize in East Africa. Crop Prot. 2018;105:10-5. Available from: https://doi.org/10.1016/j.cropro.2017.11.003