



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2015; 3 (2): 121-123

© 2015 JEZS

Received: 08-03-2015

Accepted: 03-04-2015

Ruidar Ali ShahAgriculture Research Institute,
Tarnab, Peshawar Pakistan.**Imtiaz Ali Khan**Department of Entomology,
The University of Agriculture
Peshawar, Pakistan.

Evaluation of onion cultivars against onion thrips, *Thrips tabaci* (Lindeman) infestation on onion crop

Ruidar Ali Shah, Imtiaz Ali Khan

Abstract

The experiment to evaluate the onion cultivars against *Thrips tabaci* infestation was conducted at Agricultural Research Institute Tarnab Peshawar, Pakistan during 2011-12. Eight onion cultivars (Ambika, Swat-1, Trichmir, Barkel, Macarena, Red ball, Granada Red and Sunset) were assessed to determine the most tolerant cultivar against *T. tabaci* infestation. *T. tabaci* populations were monitored on weekly basis on onion plants under natural field condition. *T. tabaci* infested all the cultivars more or less, however highest numbers of thrips plant⁻¹ were recorded in Trichmer (10.99 thrips plant⁻¹) and lowest (5.98 thrips plant⁻¹) were recorded in Swat-1. Highest plant height (64.35 cm) was recorded in cultivar Trichmir and lowest (38.40 cm) was recorded in Macarena. Highest (13.07) number of leaves plant⁻¹ was recorded in Trichmir and lowest (7.66 leaves plant⁻¹) was recorded in Granada red. Trichmir was (94.04%) the most succulent cultivar followed by Sunset (91.38%), Ambika (87.21%) and (86.49%) in cultivar Swat-1. The maximum yield (18.56 tons ha⁻¹) was obtained from cultivar Swat-1 and minimum was from Ambika (9.34 tons ha⁻¹). Number of leaves, plant height, high succulency and color favored high infestation.

Keywords: Cultivar, Peshawar, *Thrips tabaci*, yield

1. Introduction

Onion, *Allium cepa* L., belongs to the family Amaryllidaceae (Liliaceae) and is a major horticultural cash crop in Pakistan^[7]. The onion consumed in the fresh form as well as used in a frozen and dehydrated bulbs by the low and high income community in Indo-Pakistan^[2]. In Pakistan the yield of onion is low as compared to the other developing countries. There are many reasons for the low yield, among which insect pests are the important one. There are many insect pests of onion crop including thrips, maggots, head borer, cutworm and leaf minors which cause reduction in yield. Among these, onion thrips, *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) is a major pest of onion crop in Pakistan^[11]. The feeding of *T. tabaci* causes direct damage to plant while indirectly harms the crop by transmitting viruses^[16]. The nymphs do more damage than adults in a peculiar feeding behavior in flower and even in fruits^[8]. *T. tabaci* feeds in a piercing-sucking manner, leaving silvery areas on leaves, flowers and fruits and consequently consume the cell substances via the feeding tube formed by the maxillary stylets^[15, 3]. *T. tabaci* select new plant tissue for feeding and consequently the silvery patches or streaks are formed which can be easily seen during sun shine. In severe attack of *T. tabaci* the size of the onion bulb become reduced and in some cases up to 66% of the onion crop may be lost^[13]. The objective of this study was to evaluate and select the most tolerant onion cultivar commercially grown in Khyber Pakhtunkhwa Province, Pakistan against *T. tabaci* infestation.

2. Material and methods

Evaluation of eight commercially grown onion cultivars (Ambika, Swat-1, Trichmir, Barkel, Macaria, Red ball, Granada red and Sunset) against onion thrips infestation was conducted at the Agricultural Research Institute Tarnab, Peshawar, Pakistan during 2011-12.

2.1 Cultivation of crop

Healthy seedlings 6" in length approximately of the selected onion cultivars of onion were transplanted in different plots, each plot measuring 3 x 5 m². Plant to plant and row to row distance were kept at 15 cm and 30 cm respectively. Standard agronomic practices were carried out throughout the growing season of the crop for both the years.

Correspondence:**Ruidar Ali Shah**Agriculture Research Institute,
Tarnab, Peshawar Pakistan.

2.2 Data collection

Data on thrips population were recorded weekly from fifteen randomly selected plants in each treatment from each replication till maturity of the crops. Plant morphological characters like color, height, number of leaves and succulency were recorded. For these, five randomly selected mature leaves were cut and carried back to the Laboratory. Color, height and the number of leaves were recorded for each treatment. For succulency plants were selected from one square meter area and took equal weight of each cultivar for recording the fresh and dry foliage weight^[10]. After receiving the fresh weight, the vegetative part of each cultivar were marked and placed in an electric oven at 70±1 °C for 48 hours to dry up. The percent plant succulency was determined with the following formula; % Plant Succulency = 100 - (Dry Foliage weight / Fresh Foliage weight x 100)

When 70% of the foliage was dropped to the ground, the onions were dug out and the bulb yield was determined for each plot. Based on these yields, estimated yield /ha were calculated for each cultivar and were compared with other cultivars for thrips damage.

2.3 Statistical analysis of data

The experiments were laid out in Randomized Complete Block (RCB) design with three replications. Data were analyzed using the Microcomputer statistical program for experiments. ANOVA were constructed to test the significance difference between the variables. The least significant difference (LSD) tests were applied to differentiate the means.

3. Results and discussion

Thrips tabaci were observed regularly and the numbers of thrips plant⁻¹ were recorded on each cultivar until the crops were harvested. Mean numbers of thrips were significantly more (10.99 thrips plant⁻¹) in Trichmer and lowest (5.98 thrips plant⁻¹) in Swat-1, Ambika recorded significantly more number of thrips from Macarena, Red ball, Granada red and Sunset (Table 1). Our results are comparable with the findings of Alimousavi, Diaz-Montano, Sepahvand and Yousefi^[1, 4, 14 & 17] who found significant difference on number of thrips per plant between the various cultivars.

Table 1: Mean numbers of *T. tabaci* per plant on eight onion cultivars at ARI, Tarnab Peshawar, Pakistan during 2011-12.

Cultivars	Years		Means
	2011	2012	
Ambika	8.58 b	9.85 b	9.21 b
Swat-1	6.77 c	5.18 e	5.97 d
Trichmer	9.92 a	12.07 a	10.99 a
Barkel	7.77 bc	10.73 b	9.25 b
Macarena	7.87 bc	8.63 c	8.25 c
Red ball	8.39 b	7.68 cd	8.04 c
Granada Red	8.78 b	7.25 d	8.02 c
Sunset	7.90 bc	7.29 d	7.60 c
LSD	1.022	1.135	0.729

Means followed by similar letters within each column are not significantly different at 5% level of probability.

Results in the (Table 2) showed plant morphological characters observed in all the selected cultivars. Ambika,

Swat-1, Barkel and Red ball were green in color while Trichmer, Macarena, Granada red and Sunset were dark green in color. Thrips prefer dark color and those cultivars having dark in color are more susceptible to thrips infestation as reported by Ellis,^[5] Sepahvand,^[14] also observed a positive correlation in the number of thrips per plant with the foliage color. Trichmer cultivar was tallest having (64.35cm) followed by Swat-1 (53.57cm), while Macarena was the shortest (38.40 cm). However there were no significant different among the Barkel, Macarena, Red ball, Granada red and Sunset. Trichmer cultivar was taller amongst all the cultivars therefore, this cultivars was susceptible to have more infestation of *T. tabaci*, similar observation was made by Kibanyu,^[9] that taller varieties have more chances of infestation than shorter one. Trichmer, Ambika, Barkel and Swat-1 have significantly more number of leaves than Macarena, Red ball, Granada red and Sunset, however highest number of leaf (13.07) per plant were recorded in Trichmer and lowest (7.66) were recorded in Granada red. Highly leafy onion varieties have a tendency to bear a higher number of thrips as compared to less number of leaves Malik,^[10] Trichmer was the most succulent (94.04%) among the cultivars followed by Sunset (91.38%), Granada red (90.4%) and Macarena (90.13%). Swat-1 showed least succulent (86.49%) among the other cultivars. Natarajan,^[12] revealed that plant succulency has a positive impact on densities of thrips as well as other sucking insect pests.

Table 2: Plant morphological characters identified in eight onion cultivars at ARI, Tarnab Peshawar, Pakistan during 2011-12.

Cultivars	Plant color	Plant Height (cm)	Number of leaves/plant	Succulency (%)
Ambika	Green	41.80 c	12.90 a	87.21 c
Swat-1	Green	53.57 b	11.48 a	86.49 c
Trichmir	Dark Green	64.35 a	13.07 a	94.04 a
Barkel	Green	42.28 c	11.91 a	87.21 c
Macarena	Dark Green	38.40 c	8.66 b	90.13 b
Red ball	Green	39.62 c	8.53 b	89.01 bc
Granada Red	Dark Green	41.98 c	7.66 b	90.94 b
Sunset	Dark Green	38.50 c	8.07 b	91.38 ab
LSD		4.31	1.558	2.77

Means followed by similar letters within each column are not significantly different at 5% level of probability.

Highest yield (18.56 tons ha⁻¹) was recorded in the cultivar Swat-1 and lowest (9.34 tons ha⁻¹) in Ambika (Table 3). Cultivar Swat-1 proved to be the most tolerant against onion thrips infestation and gave high yield in comparison to other cultivars. Similar finding was recorded by Sepahvand,^[14] they found significant difference between the cultivars in relation to yield, duration of maturity and number of thrips plant⁻¹. Fournier,^[6] also observed reduction on yield on the plant which were high numbers of thrips infestation. On the basis of these findings it is concluded that cultivar Swat-1 proved to be the most tolerant against thrips infestation and gave high yield in comparison to other cultivars.

Table 3: Yield tons ha⁻¹ in each cultivar at ARI, Tarnab Peshawar, Pakistan during 2011-12.

Cultivars	Years		Means
	2011	2012	
Ambika	8.92 c	9.77 c	9.34 e
Swat-1	17.76 a	19.35 a	18.56 a
Trichmer	13.05 b	10.61 c	11.83 d
Barkel	12.37b	15.43 b	13.90 cd
Macarena	14.75 b	16.81 ab	15.78 bc
Red ball	13.77 b	15.93 b	14.85 bc
Granada Red	14.58 b	16.78ab	15.68 bc
Sunset	14.72 b	17.73ab	16.23 b
LSD	2.97	3.32	2.13

Means followed by similar letters within each column are not significantly different at 5% level of probability.

4. Acknowledgement

This study was supported by Higher Education Commission of Pakistan (HEC) and is a part of Ph.D. research work. The authors are also grateful to the Entomologist, field staff of Entomology section and Director General Agricultural Research Institute Tarnab for providing space in their experimental field.

5. References

- Alimousavi SA, Hassandokht MR, Moharramipour S. Evaluation of Iranian onion germplasms for resistance to thrips. *International Journal of Agriculture and Bioscience* 2007; 9:897-900.
- Baloch A. Vegetable crops. *Horticulture*, 1994, 525-526.
- Chisholm I, Lewis T. A new look at thrips (Thysanoptera) mouthparts, their action and effects of feeding on plant tissue. *Bulletin of Entomological Research* 1984; 74(04):663-675.
- Diaz-Montano J, Fuchs M, Nault BA, Shelton AM. Evaluation of onion cultivars for resistance to onion thrips, *Thrips tabaci* (Thysanoptera: Thripidae) and Iris yellow spot virus. *Journal Economic Entomology* 2010; 103(3):925-937.
- Ellis BW, Bradley FM, Atthowe H. *The Organic Gardener's Handbook of Natural Insect and Disease Control: A Complete Problem-Solving Guide to Keeping Your Garden and Yard Healthy without Chemicals*: Rodale, 1996.
- Fournier F, Boivin G, Stewart RK. Effect of *Thrips tabaci* (Thysanoptera: Thripidae) on yellow onion yields and economic thresholds for its management. *Journal of Economic Entomology* 1995; 88(5):1401-1407.
- Hassan S, Malik MF. Weeds management in broadcasted onion (*Allium cepa*). *Asian Journal of Plant Science* 2002; 1(1):28-30.
- Kawai A. Studies on population ecology of *Thrips palmi* Karny. 16. Distribution among leaves, flowers and fruit on aubergine and sweet pepper. *Japanese Journal of Applied Entomology and Zoology* 1988; 32(4):291-296.
- Kibanyu JK. Survey of Production Practices and Evaluation of Onion Varieties Susceptibility to Thrips in Kirinyaga District, Kenya, 2013.
- Malik MF, Nawaz M, Hafeez Z. Evaluation of promising onion (*Allium cepa*) varieties against thrips infestation on the agro-ecosystem of Balochistan, Pakistan. *Asian Journal of Plant Science* 2003; 2:716-718.
- Malik MF, Rashid M, Iqbal J, Ahmad A. Resistance determination against thrips of promising onion varieties in the agro-ecosystem of Balochistan, Pakistan. *Punjab University Journal of Zool* 2010; 25(1-2):1-11.
- Natarajan K. Influence of NPK fertilization on the population density of cotton whitefly. Paper presented at: Proceedings of National Symposium of Resurgence of sucking pests, 1986.
- Ochoa MDL, Zavaleta-Mejía E, Johansen N, Herrera G, Soriano EC. Tospoviruses, weeds and thrips associated with chrysanthemum (*Dendranthema grandiflora* Tzvelev cv. Polaris). *International Journal of Pest Management* 1996; 42(3):157-159.
- Sepahvand N, Jaliyani N, Abbasi Far AR. Evaluation of Resistance and Susceptible Iranian Onion Cultivars and Landraces to Thrips in Karaj and Arak. *Agriculture Research Educational Organization* 2009, 38.
- Stafford CA, Walker GP, Ullman DE. Infection with a plant virus modifies vector feeding behavior. *Proceedings of the National Academy of Sciences* 2011; 108(23):9350-9355.
- Whitfield AE, Ullman DE, German TL. Tospovirus-thrips interactions. *Annual Review of Phytopathol* 2005; 43:459-489.
- Yousefi M, Abasifar A, Hafshejani AF, Sendi JJ. Resistance of eight Iranian onion cultivars to onion thrips (*Thrips tabaci* Lindeman) in the Markazi Province of Iran. *African Journal of Agriculture Research* 2011; 6(21):4925-4930.