



E-ISSN: 2320-7078  
P-ISSN: 2349-6800  
JEZS 2015; 3(5): 235-238  
© 2015 JEZS  
Received: 14-08-2015  
Accepted: 15-09-2015

**Anita Singh**  
Division of Entomology,  
Department of Zoology, Faculty  
of Science, The Maharaja  
Sayajirao University of Baroda,  
Vadodara-390002

**Dolly Kumar**  
Division of Entomology,  
Department of Zoology, Faculty  
of Science, The Maharaja  
Sayajirao University of Baroda,  
Vadodara-390002

**Correspondence:**  
**Dolly Kumar**  
Division of Entomology,  
Department of Zoology, Faculty  
of Science, The Maharaja  
Sayajirao University of Baroda,  
Vadodara-390002

## Feeding potential of *Scymnobiussordidus* Horn (Family: Coccinellidae) on cotton mealybug, *Phenacoccus solenopsis* (Tinsley)

**Anita Singh, Dolly Kumar**

### Abstract

During 2008-2011 five natural enemies against mealybug *Phenacoccus solenopsis* were recorded from agriculture fields of Vadodara. Out of these natural enemies *Scymnobiussordidus* was used for recording feeding potential against mealybug. The result shows that the fourth instar larvae were voracious feeder. It was feeding around  $36.2 \pm 0.42$  number of mealybug nymphs and  $2.8 \pm 0.2$  number of mealybug adults. In its life cycle, the larva stages consumed around  $61.6 \pm 0.48$  nymphs and  $7.6 \pm 0.48$  adults of mealybug. Hence, *S. sordidus* is having good potential as bio control agents against mealybug.

**Keywords:** *Phenacoccus solenopsis*, *Scymnobiussordidus*, Natural enemies

### 1. Introduction

Today management of mealybug *P. solenopsis* is major concern in the agricultural fields. *P. solenopsis* is the major menace in many tropical and subtropical countries; attacking 159 hosts plant species (agricultural and horticultural crops) belonging to 21 different families [24, 31]. In India this *P. solenopsis* is taken prime position in North India (Punjab, Haryana and Rajasthan), whereas it shows moderate incidence of damage in Central India (Gujarat and Maharashtra) and South India [9]. It cause yield loss by direct and indirect infestation. In direct mode of infestation they feed on leaves, stems, fruits and roots of host plants. Whereas in indirect mode of infestation they secrete honey dew which causes development of sooty mould which inhibit photosynthesis which leads to host plant death [23]. Due to this direct and indirect infestation by mealybug causes huge yield loss; reducing the cotton yield by up to 40-50 % in infested fields [16]. This yield loss leads the grower to protect their crops by applying synthetic chemical insecticides such as organophosphates, carbonates, synthetic pyrethroids and nicotinoids [17]. Although indiscriminate use of these chemicals are showing direct impacts on natural enemies life span, fecundity and ability to locates hosts which encourages the use of biorational alternatives for minimizes the load of pesticides [6, 1, 10, 21]. Therefore present study was conducted for identification of various natural enemies against mealybugs. Further the consumption rate of one of the predator, *Scymnobiussordidus* Horn was recorded in laboratory to know its feeding potential so that it can be used in future.

### 2. Materials and methods

Observations for naturally found bio-control agents associated with mealybugs were recorded from the selected agricultural sites in Vadodara from 2008 to 2011 at weekly interval from 25 randomly selected plants having infestation.

#### Collection, preservation and identification of bio control agents associated with mealybug

Each of the study sites was visited twice a month. On each day the sampling was done twice once in morning hrs (7 am to 9 am) and second time during evening hrs (5 pm to 7 pm). Collection was done by sweeping and hand picking methods. Insects were killed using killing jars with chloroform. Collected insects are pinned accordingly. Collected larval stages were transferred in to vials having 70-90% ethyl alcohol, brought to the laboratory, mounted on slides and then observed and identified. Identification and labeling was done using keys available in standard taxonomic literature [2, 5].

**Mass multiplication of *Phenacoccus solenopsis*** <sup>[20]</sup>

The rearing and breeding of *P. solenopsis* was carried at room temperature (24.3-31.2 °C, 48.9-71.7% R.H.) in the laboratory. The cotton twigs, harboring *P. solenopsis* colonies, were collected from the fields and brought to the laboratory. The mealybugs present on these cotton twigs were gently removed with the help of soft camel hair brush and were released on the cotton twigs (Twigs end was wrap with wet cotton) kept in plastic box (20 x 15 cm<sup>2</sup>) in laboratory. Newly hatched crawlers were collected from the ovipositing female of laboratory culture and were placed onto the cotton twigs for mass rearing. Culture obtained was used for further experiments.

**Mass Multiplication of *Scymnobijs sordidus***

A predator stock culture was obtained from *P. solenopsis* collected from cotton and okra fields in Vadodara. The predatory larva was reared on *P. solenopsis* grown on hibiscus leaves and twigs in plastic cages as described above for rearing of mealybugs. 5 pairs of male and female were transferred into a small plastic cage containing 3 instars and virgin females of mealybug. Culture was maintained at 28 ± 1 °C, 60 ± 5% RH, under a photoperiod of 16:8 L: D.

**To assess the feeding potential**

After hatching 25 nymphs and adults of *P. solenopsis* were offered to 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> larval stage of *S. sordidus* separately. 10 insects were added daily after 1<sup>st</sup> day. Single larva of *S. sordidus* was released after one hour of release of nymphs and adults in each petri plate (after settling of mealybugs). The predatory potential was recorded by counting the number of mealybugs fed by 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> larval stage of *S. sordidus* up to pupation.

**3. Results and Discussion**

The survey to the study sites shows that 5 species of biological control agents have been recorded feeding on cotton mealybugs in the agricultural fields of Vadodara (Figure 1A, 1B, 1C, 1D, & 1E).

Figure 1A, 1B, 1C, 1D, 1E: Natural enemies collected from Vadodara agriculture field



1A. Larvae of *Scymnobijs sordidus* Horn



1B. *Aenasijs bambawalei* Hayat & its cocoons



1C. *Sympherobius fallax* (Navas)



1D. *Chrysoperla zastrowi Arabica* (Henry)



1E. *Cheilomenes sexmaculata* Fab.

These field collected natural enemies was from three different orders (coleoptera, Neuroptera, hymenoptera) and 4 different families (Table 1).

**Table 1:** Biocontrol agents recorded for *Phenacoccus solenopsis* from agriculture field of Vadodara

Order	Family	Scientific name	Status
Neuroptera	Hermerobiidae	<i>Sympherobius fallax</i> (Navas)	Predator
	Chrysopidae	<i>Chrysoperla zastrowi Arabica</i> (Henry)	
Coleoptera	Coccinellidae	<i>Cheilomenes sexmaculata</i> (Fab.)	
		<i>Scymnobijs sordidus</i> (Horn)	
Hymenoptera	Encyrtidae	<i>Aenasijs bambawalei</i> Hayat	Parasitoid

Similarly, seven species of predatory coccinellid beetles belonging to sub family Scymninae from vegetable field of Kerala associated with sucking pest was reported <sup>[12]</sup>. All collected species are having great potential against mealybugs. *Aenasijs bambawalei* is reported as good natural enemy of mealybug *P. solenopsis* in Gujarat and Haryana <sup>[13, 11]</sup>. It is having high parasitism potential against invasive mealybug *P.*

*solenopsis* [7]. Predators such as lacewing *Chrysoperla carnea* and *Symphorobius fallax* are having high capability for controlling mealybug *P. solenopsis* [18, 8, 15]. Although during field visit it was also recorded that the farmers were driven to extensive usage of insecticides against mealybug in Vadodara [22]. These practices were showing pest resurgence, secondary pest outbreak and adverse effect on natural enemies (non-target effects) of the pest. The adverse effect of broad spectrum insecticides like chlorpyrifos caused adverse effect on *Coccidoxenoides peregrinus* (Timberlake) (Hymenoptera: Encyrtidae) of *Planococcus ficus* (Signoret) [25]. Similarly in South Africa use of contact insecticides like synthetic pyrethroid (Cypermethrin), Carbamate (Mancozeb) against Vine mealybugs showed adverse effect on the development of parasitoids *Anagyrus sp.* and *Coccidoxenoides perminutus* [14]. These negative impacts on non target insects make chemical control as less desirable option and insist on development of proper strategies to control the *P. solenopsis* by slowly weaning away the farmers from use of pesticides to a more ecofriendly pest management like use of bio-control agents. Out of all identified natural enemies the *Scymnobioides sordidus* Horn was used for evaluation of its feeding potential. The result shows that the fourth instar larvae were voracious feeder. This stage was feeding around  $36.2 \pm 0.42$ ,  $2.8 \pm 0.2$  number of mealybug nymphs and adults respectively. Larvae were mainly preferred nymph than adult of mealybug. The total number of *P. solenopsis* nymph, adult consumed by the larvae of *S. sordidus* during its life cycle was  $61.6 \pm 0.48$  and  $7.6 \pm 0.48$  respectively (Table 2).

**Table 2:** Consumption of different stages of *Phenacoccus solenopsis* Tinsley by different instars of *Scymnobioides sordidus*

Mealybug Stages	<i>Scymnobioides sordidus</i> Larval Instars (Mean $\pm$ SE)				
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	Total
Nymphs	5.8 $\pm$ 0.13	7.2 $\pm$ 0.2	12.4 $\pm$ 0.16	36.2 $\pm$ 0.42	61.6 $\pm$ 0.48
Adults	1.4 $\pm$ 0.16	1.6 $\pm$ 0.16	1.8 $\pm$ 0.13	2.8 $\pm$ 0.2	7.6 $\pm$ 0.48

Similar result was reported by many Coccinellid species such as *Cryptolaemus montrouzieri*, *Brumus suturalis* which preferred nymph stage than adult of *Phenacoccus solenopsis* [4, 10, 18, 19]. The high potential of predatory nature gives good indication to use *Scymnobioides sordidus* Horn as bio-control agent for *Phenacoccus solenopsis*.

#### 4. Acknowledgement

One of the authors (Anita Singh) sincerely thanks UGC for grants provided under RFSMS scheme.

#### 5. References

- Ambika, Selvisabhanayakam. Impact of vijay neem on histological and scanning electron microscopic changes in the testis of adult male insect *Odobtopus vericornis* (Dist) (Heteroptera: Pyrrhocoridae). International Journal of Pharma and Bio Sciences. 2012; 3:460-466.
- Ananthkrishnan TN, David BV. (2004) General and applied entomology. Tata McGraw Hill publishing company limited, New Delhi.
- Arif MI, Rafiq M, Ghaffar A. Host plant of cotton mealybug (*Phenacoccus solenopsis*): a new menace to cotton agroecosystem of Punjab. International Journal of Agriculture and Biology. 2009; 11:163-167.
- Atlihan R, Kaydan B, Ozgokee MS. Feeding activity and life history characteristics of generalist predator, *Chrysoperla carnea* (Neuroptera: Chrysopidae) at different prey densities. Journal of pest Science. 2004; 77:17-21.
- Borror DJ, Triplehorn CA, Johnson NF. An introduction to study of insects. Sixth edition. Saunders college publishing, 1992.
- Cork A, Kamal NQ, Alam SN, Chaudhary CJS, Talekar NS. Pheromones and their applications to insect pest control – A review. Bangladesh Journal of Entomology. 2003; 13:1-13.
- Feng D, Li P, Zhou Z, Xu Z. Parasitism potential of *Aenasius bambawalei* on the invasive mealybug *Phenacoccus solenopsis*. Biocontrol Science and Technology 2014; 24:1333-1338.
- Gillani WA, Copland M, Raja S. Studies on feeding preference on brown lacewing (*Symphorobius fallax* Navas) larvae for different stages of long tailed mealybug (*Pseudococcus longispinus*) (Targioni-Tozzetti). Pakistan Entomologist 2009; 31:1-4.
- Hanchinal SG, Patil BV, Bheemanna M, Hosamani AC. Population dynamics of mealybug, *Phenacoccus solenopsis* Tinsley and its natural enemies on Bt cotton. Karnataka Journal of Agricultural Sciences. 2010; 23:137-139.
- Harmeet K, Virk JS. Feeding potential of *Cryptolaemus montrouzieri* against the mealybug *Phenacoccus solenopsis*. Phytoparasitica 2012; 40:131-136.
- Jhala RC, Solanki RF, Bharpoda TM, Patel MG. Occurrence of Hymenopterous Parasitoids, *Aenasius bambawalei* Hayat and *Promuscidea unfaciativentris* Girault on Cotton Mealy Bugs, *Phenacoccus solenopsis* Tinsley in Gujarat. Insect Environment 2009, 14:164-165.
- Jose Q, Bhaskar H, Mathew MP. Predaceous symninae (Coccinellidae: Coleoptera) associated with sucking pests of vegetables. Journal of insect science 2011; 24:13-17.
- Kumar R, Kranthi KR, Monga D, Jat SL. Natural parasitization of *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) on cotton by *Aenasius bambawalei* Hayat (Hymenoptera: Encyrtidae). Journal of Biological Control. 2009; 23(4):457-460.
- Mgocheki N, Addison P. Effect of contact pesticides on vine mealybug parasitoids, *Anagyrus sp.* near *Pseudococci* (Girault) and *Coccidoxenoides perminutus* (Timberlake) (Hymenoptera: Encyrtidae). South African Journal of Entology and Viticulture. 2009, 30:110-116.
- Muzammil S, Muhammad H, Sajid N. Predatory potential of *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae) against cotton mealybug. Pakistan Entomologist 2007; 29:103-106.
- Nagrare VS, Kranthi S, Biradar VK, Zade NN, Sangode V, Kakde G *et al.* Widespread infestation of the exotic mealybug species *Phenacoccus solenopsis* (Tinsley) (Hemiptera: Pseudococcidae) on cotton in India. Bulletin of Entomological Research 2009; 99:537-41.
- Nas MN. *In vitro* studies on some natural beverages as botanical pesticides against *Erwinia amylovora* and *Curobacterium flaccumfaciensis* subsp. poinsettiae. Turkish Journal of Agriculture and Forestry. 2004; 28:57-61.
- Rashid MMU, Khattak MK, Abdullah K, Amir M, Tariq M, Nawaz S. Feeding potential of *Chrysoperla carnea* and *Cryptolaemus montrouzieri* on cotton mealybug, *Phenacoccus solenopsis*. The Journal of Animal and Plant Sciences. 2012, 22: 639-643.
- Shah N, Mohammad K, Ghulam HA, Muzaffar AT, Rabdino K. Feeding potential of lady bird beetle, *Brumus suturalis* (Coleoptera: Coccinellidae) on cotton mealybug *Phenacoccus solenopsis* (Tinsley) in laboratory and field. Sarhad Journal of Agriculture. 2012; 28:259-265.

20. Singh A, Kumar D. Population dynamics, biology of mealybug *Phenacoccus solenopsis* (Tinsley) and its natural enemies in Vadodara, Gujarat. Recent Research in Science and Technology. 2012, 4:22-27.
21. Singh A, Kumar D. Studies on presence of semiochemicals from *Phenacoccus solenopsis* (Tinsley). International Journal of Pharma and Bio Sciences. 2015; 6:326-332.
22. Singh A, Kataria R, Kumar D. Repellence property of traditional plant leaf extracts against *Aphis gossypii* Glover and *Phenacoccus solenopsis* Tinsley. African Journal of Agricultural Research. 2012; 7:1623-1628.
23. Vennila S, Ramamurthy VV, Deshmukh A, Pinjarkar DB, Agarwal M, Pagar PC *et al.* A treatise on mealybugs of central Indian cotton production system, Technical Bulletin no. 24, National Centre for Integrated Pest Management, New Delhi, 2010, 39.
24. Wang Y, Watson GW, Zhang R. The potential distribution of an invasive mealybug *Phenacoccus solenopsis* and its threat to cotton in Asia. Agricultural and Forest Entomology 2010; 12:403-41.
25. Walton VM, Pringle KL. Effect of pesticides used on table grapes on the mealybug parasitoids *Coccidoxenoides peregrinus* (Timberlake) (Hymenoptera: Encyrtidae). South African Journal of Enology and viticulture. 1990, 20:31-34.